







University of the State of New York

---

BULLETIN

OF THE

New York State Museum

FREDERICK J. H. MERRILL *Director*

No. 33 Vol. 7

April 1900

---

CHECK LIST

OF

NEW YORK BIRDS

By

MARCUS S. FARR D.Sc.


ALBANY

UNIVERSITY OF THE STATE OF NEW YORK

1900

M65m-Fo-1500

Price 25 cents



# University of the State of New York

## REGENTS

With years of election

1874 ANSON JUDD UPSON L.H.D. D.D. LL.D.

*Chancellor, Glens Falls*

1892 WILLIAM CROSWELL DOANE D.D. LL.D.

*Vice-Chancellor, Albany*

1873 MARTIN I. TOWNSEND M.A. LL.D. - - - Troy

1877 CHAUNCEY M. DEPEW LL.D. - - - New York

1877 CHARLES E. FITCH LL.B. M.A. L.H.D. - - Rochester

1877 ORRIS H. WARREN D.D. - - - Syracuse

1878 WHITELAW REID LL.D. - - - New York

1881 WILLIAM H. WATSON M.A. M.D. - - - Utica

1881 HENRY E. TURNER - - - Lowville

1883 ST CLAIR MCKELWAY L.H.D. LL.D. D.C.L. - - Brooklyn

1885 HAMILTON HARRIS Ph.D. LL.D. - - - Albany

1885 DANIEL BEACH Ph.D. LL.D. - - - Watkins

1888 CARROLL E. SMITH LL.D. - - - Syracuse

1890 PLINY T. SEXTON LL.D. - - - Palmyra

1890 T. GUILFORD SMITH M.A. LL.D. C.E. - - - Buffalo

1893 LEWIS A. STIMSON B.A. M.D. - - - New York

1895 ALBERT VANDER VEER Ph.D. M.D. - - - Albany

1895 CHARLES R. SKINNER M.A. LL.D.

Superintendent of Public Instruction, ex officio

1897 CHESTER S. LORD M.A. LL.D. - - - Brooklyn

1897 TIMOTHY L. WOODRUFF M.A. Lieutenant-Governor, ex officio

1899 THEODORE ROOSEVELT B.A. LL.D. Governor, ex officio

1899 JOHN T. McDONOUGH LL.B. LL.D. Secretary of State, ex officio

1900 THOMAS A. HENDRICK M. A. - - - Rochester

## SECRETARY

Elected by regents

1900 JAMES RUSSELL PARSONS JR M.A.

## DIRECTORS OF DEPARTMENTS

1888 MELVIL DEWEY M.A. *State library and Home education*

1890 JAMES RUSSELL PARSONS JR M.A.

*Administrative, College and High school dep'ts*

1890 FREDERICK J. H. MERRILL Ph.D. *State museum*



200  
52  
University of the State of New York

---

BULLETIN  
OF THE  
New York State Museum

FREDERICK J. H. MERRILL *Director*

No. 33 Vol. 7

April 1900

---

CHECK LIST  
OF  
NEW YORK BIRDS

By  
MARCUS S. FARR D. Sc.

ALBANY  
UNIVERSITY OF THE STATE OF NEW YORK

1900



## PREFACE

This list was originally prepared for use in collecting data for a bulletin on the birds of New York, but, inasmuch as no check list of the birds of the entire state is available for the student of our avi-fauna, it has been thought advisable to publish it in this form, hoping that it may be of some use to students in ornithology and serve as a convenient means of keeping notes on the various species.

My purpose has been to include all the birds known to occur or to have occurred in our state, and only those forms have been admitted that have actually been taken in New York. Thus the list contains not only our regular summer and winter residents and migrants, but also those that occur as irregular visitants or stragglers. A number of the species recorded have been taken only once or twice in the state and this is in nearly every case indicated in a foot note with the proper reference. It is believed that nearly, if not quite all the birds of our state, that have been recorded, are included; but for any corrections or additions the author will be grateful.

The list is divided into two parts, the first containing those birds that have actually been taken in New York, the second including those of which we have no satisfactory state records, but which, having been taken in adjoining states, may be looked for in New York.

Strict adherence is given to the nomenclature of the American ornithologists union check list of 1895 and the supplements of January 1897 and 1899. The number of the species in this list is followed by the American ornithologists union number in parentheses or brackets, and as the pamphlet is designed for the use of students rather than advanced workers, the common name is given first.


My obligations are due to Dr F. J. H. Merrill, whose kind aid and criticism have been a source of inspiration in my work. Also to Dr J. A. Allen and Mr William Dutcher of New York, Dr William Brewster of Cambridge, and Dr A. W. Butler of Indianapolis, who have been so kind as to give me notes and suggestions on several species.

MARCUS S. FARR

*Assistant zoologist*

*New York state museum*

*8 Feb. 1900*



# CHECK LIST OF NEW YORK BIRDS

---

## PART I

### Order **Pygopodes**    *Diving birds*

#### Family **PODICIPIDAE**    *Grebes*

- 1    (2) Holboell's grebe  
      *Colymbus holboellii* (Reinh.)

- 2    (3) Horned grebe  
      *Colymbus auritus* Linn.

- 3    (6) Pied-billed grebe  
      *Podilymbus podiceps* (Linn.)

#### Family **GAVIIDAE**    *Loons*

- 4    (7) Loon  
      *Gavia imber* (Gunn.)





- 5 (9) Black-throated loon<sup>1</sup>  
Gavia artica (Linn.)

- 6 (11) Red-throated loon  
Gavia lumme (Gunn.)

Family ALCIDAE    *Auks, murrees and puffins*

- 7 (13) Puffin  
Fratercula arctica (Linn.)

- 8 (27) Black guillemot  
Cepphus grylle (Linn.)

---

<sup>1</sup> One New York record, Long Island. See *Auk*, 10:265-66.



- 9 (31) Brünnich's murre  
    *Uria lomvia* (Linn.)

- 10 (32) Razor-billed auk  
    *Alca torda* Linn.

- 11 (34) Dovekie  
    *Alle alle* (Linn.)

Order **Longipennes**    *Long-winged swimmers*

Family STERCORARIIDAE    *Skuas and jaegers*

- 12 (35) Skua<sup>1</sup>  
    *Megalestris skua* (Brünn.)

---

<sup>1</sup> Long Island record, *Auk*, 3 : 432-33. Niagara river, spring of 1886. See *Auk*, 6 : 331; also Bergtold's *List of the birds of Buffalo and vicinity*, p. 3.





- 13 (36) Pomarine jaeger  
Stercorarius pomarinus (Temm.)

- 14 (37) Parasitic jaeger  
Stercorarius parasiticus (Linn.)

- 15 (38) Long-tailed jaeger  
Stercorarius longicaudus Vieill.

Family LARIDAE    *Gulls and terns*

- 16 (39) Ivory gull<sup>1</sup>  
Pagophila alba (Gunn.)

---

<sup>1</sup> One Long Island record. See *Auk*, 12: 290-91.



- 17 (40) Kittiwake  
*Rissa tridactyla* (Linn.)

- 18 (42) Glaucous gull  
*Larus glaucus* Brünn.

- 19 (43) Iceland gull  
*Larus leucopterus* Faber

- 20 (47) Great black-backed gull  
*Larus marinus* Linn.



21 (51) Herring gull<sup>1</sup>*Larus argentatus* Brünn.

## 22 (51a) American herring gull

*Larus argentatus smithsonianus* Coues

## 23 (54) Ring-billed gull

*Larus delawarensis* Ord

## 24 (58) Laughing gull

*Larus atricilla* Linn.

---

<sup>1</sup> Dr J. A. Allen writes me that this species is occasional in winter in Long Island sound and adjacent waters, and that several specimens taken in that locality and referable to this form have come under his observation.





- 25 (60) Bonaparte's gull  
Larus philadelphia (Ord)

- 26 [60.1] Little gull<sup>1</sup>  
Larus minutus Pall.

- 27 (62) Sabine's gull<sup>2</sup>  
Xema sabinii (Sab.)

- 28 (63) Gull-billed tern  
Gelochelidon nilotica (Hasselq.)

---

<sup>1</sup> There is one American record of the occurrence of this European species. It was taken at Fire Island, Long Island. See *Auk*, 1888, 5: 172.

<sup>2</sup> An arctic form. Two Long Island records. See Giraud's *Birds of Long Island*, 1844, p. 363; Baird, Brewer & Ridgway's *Water birds*, 2:272; *Auk*, 1900, 17: 63.



- 29 (64) Caspian tern  
Sterna caspia Pallas

- 30 (65) Royal tern  
Sterna maxima Bodd.

- 31 [68] Trudeau's tern<sup>1</sup>  
Sterna trudeaui Aud.

- 32 (69) Forster's tern  
Sterna forsteri Nutt.

---

<sup>1</sup> One representative of this South American bird has been recorded from Long Island. See Aud. *Orn. biog.* 5:125; Aud. *Syn. birds N. A.*, p. 319; Giraud's *Birds of Long Island*, p. 354; Baird, Brewer & Ridgway's *Water birds*, 2:291.





## 33 (70) Common tern

*Sterna hirundo* Linn

## 34 (71) Arctic tern

*Sterna paradisaea* Brünn.

## 35 (72) Roseate tern

*Sterna dougalli* Montag.

## 36 (74) Least tern

*Sterna antillarum* (Less.)



- 37 (75) Sooty tern  
Sterna fuliginosa Gmel.

- 38 (77) Black tern  
Hydrochelidon nigra surinamensis (Gmel.)

Family RYNCHOPIDAE    *Skimmers*

- 39 (80) Black skimmer  
Rynchops nigra Linn.

Order **Tubinares**    *Tube-nosed swimmers*

Family PROCELLARIIDAE    *Fulmars and shearwaters*

- 40 (88) Cory's shearwater  
Puffinus borealis Cory



- 41 (89) Greater shearwater  
*Puffinus gravis* (O'Reilly)

- 42 (92) Audubon's shearwater  
*Puffinus auduboni* Finsch

- 43 (94) Sooty shearwater  
*Puffinus fuliginosus* Strickland

- 44 [98] Black-capped petrel<sup>1</sup>  
*Aestrelata hasitata* (Kuhl)

---

<sup>1</sup> Long Island record, *Ann. lyc. nat. hist. N. Y.*, 5: 220. Also Oneida lake and Ulster co. records, *Auk*, 11: 162-63; 12: 179.



45 [99] Scaled petrel<sup>1</sup>*Aestrelata scalaris* Brewst.

## 46 (106) Leach's petrel

*Oceanodroma leucorhoa* (Vieill.)

## 47 (109) Wilson's petrel

*Oceanites oceanicus* (Kuhl)Order **Steganopodes**    *Totipalmate swimmers*Family PHAËTHONTIDAE    *Tropic birds*48 (112) Yellow-billed tropic bird<sup>2</sup>*Phaëthon americanus* Grant

---

<sup>1</sup> The single specimen taken in Livingston co. in 1880 is unique, as it has not been taken elsewhere. See *Bul. Nutt. orn. club*, 6 : 91-97; *Auk*, 11 : 389-93.

<sup>2</sup> This tropical form has once been taken at Knowlesville, Orleans co. See *Bul. Nutt. orn. club*, 5 : 63. See also Short's *Birds of western New York*. Also Langille's, *Our birds in their haunts*, p. 615. One other state record. See Maynard's *Birds of eastern North America*, 1881, p. 474





Family SULIDAE    *Gannets*

- 49 (115) Booby<sup>1</sup>  
Sula sula (Linn.)

- 50 (117) Gannet  
Sula bassana (Linn.)

Family PHALACROCORACIDAE    *Cormorants*

- 51 (119) Cormorant  
Phalacrocorax carbo (Linn.)

- 52 (120) Double-crested cormorant  
Phalacrocorax dilophus (Swain.)

---

<sup>1</sup> Accidental on Moriches bay, Long Island. See *Auk*, 10: 271.



Family PELECANIDAE *Pelicans*

- 53<sub>1</sub> (125) American white pelican  
*Pelecanus erythrorhynchos* Gmel.

Family FREGATIDAE *Man-o'-war birds*

- 54 (128) Man-o'-war bird<sup>1</sup>  
*Fregata aquila* Linn.

Order **Anseres** *Lamellirostral swimmers*Family ANATIDAE *Ducks, geese and swans*

- 55<sub>1</sub> (129) American merganser  
*Merganser americanus* (Cass.)

- 56 (130) Red-breasted merganser  
*Merganser serrator* (Linn.)

---

1\_There are two Long Island records. See *American naturalist*, 9 : 470; *Auk*, 5 : 173.



57 (131) Hooded merganser  
*Lophodytes cucullatus* (Linn.)

58 (132) Mallard  
*Anas boschas* Linn.

59 (133) Black duck  
*Anas obscura* Gmel.

60 (135) Gadwall  
*Chaulelasmus streperus* (Linn.)



61 (136) Widgeon

*Mareca penelope* (Linn.)

62 (137) Baldpate

*Mareca americana* (Gmel.)

63 (139) Green-winged teal

*Nettion carolinensis* (Gmel.)

64 (140) Blue-winged teal

*Querquedula discors* (Linn.)





- 65 (142) Shoveller  
*Spatula clypeata* (Linn.)

- 66 (143) Pintail  
*Dafila acuta* (Linn.)

- 67 (144) Wood duck  
*Aix sponsa* (Linn.)

- 68 [145] Rufous-crested duck<sup>1</sup>  
*Netta rufina* (Pall.)

---

<sup>1</sup> This European species is known in the United States only by one specimen found in Fulton market, New York, and supposed to have been shot on Long Island. See *Bul. Nutt. orn. club*, 6:173; also *Proc. U. S. nat. mus.* 1881, p. 23-24.



69 (146) Redhead  
*Aythya americana* (Eyt.)

70 (147) Canvas-back  
*Aythya vallisneria* (Wils.)

71 (148) American scaup duck  
*Aythya marila* (Linn.)

72 (149) Lesser scaup duck  
*Aythya affinis* (Eyt.)



- 73 (150) Ring-necked duck  
*Aythya collaris* (Donov.)

- 74 (151) American golden-eye  
*Clangula clangula americana* Faxon

- 75 (152) Barrow's golden-eye  
*Clangula islandica* (Gmel.)

- 76 (153) Buffle-head  
*Charitonetta albeola* (Linn.)



77 (154) Old-squaw

*Harelda hyemalis* (Linn.)

78 (155) Harlequin duck

*Histrionicus histrionicus* (Linn.)

79 (156) Labrador duck<sup>1</sup>

*Camptolaimus labradorius* (Gmel.)

80 (160) American eider

*Somateria dresseri* Sharpe

---

<sup>1</sup> Formerly a winter visitant to Long Island from the arctic regions. Now supposed to be extinct.





- 81 (162) King eider  
*Somateria spectabilis* (Linn.)

- 82 (163) American scoter  
*Oidemia americana* Swains.

- 83 (165) White-winged scoter  
*Oidemia deglandi* Bonap.

- 84 (166) Surf scoter  
*Oidemia perspicillata* (Linn.)



- 85 (167) Ruddy duck  
*Erismatura jamaicensis* (Gmel.)
- 86 (169a) Greater snow goose  
*Chen hyperborea nivalis* (Forst.)
- 87 (169.1) Blue goose  
*Chen caerulescens* (Linn.)
- 88 (171a) American white-fronted goose  
*Anser albifrons gambeli* (Hartl.)



- 89 (172) Canada goose  
    *Branta canadensis* (Linn.)

- 90 (172a) Hutchins's goose  
    *Branta canadensis hutchinsii* (Rich.)

- 91 (173) Brant  
    *Branta bernicla* (Linn.)

- 92 (174) Black brant  
    *Branta nigricans* (Lawr.)



- 93 [175] **Barnacle** goose<sup>1</sup>  
    *Branta leucopsis* (Bechst.)

- 94 (180) Whistling swan  
    *Olor columbianus* (Ord)

Order **Herodiones**   *Heron, ibis, etc.*

Family IBIDIDAE   *Ibises*

- 95 (184) White ibis  
    *Guara alba* (Linn.)
- 96 (186) Glossy ibis  
    *Plegadis autumnalis* (Hasselq.)

---

<sup>1</sup> One Long Island record of this European species. See *Bul. Nutt. orn. club*, 2:18.





- 97 (187) White-faced glossy ibis<sup>1</sup>  
*Plegadis guarauna* (Linn.)

Family CICONIIDAE *Storks and wood ibises*

- 98 (188) Wood ibis  
*Tantalus loculator* Linn.

Family ARDEIDAE *Hérons, bitterns, etc.*

- 99 (190) American bittern  
*Botaurus lentiginosus* (Montag.)

- 100 (191) Least bittern  
*Ardetta exilis* (Gmel.)

---

<sup>1</sup> There is a specimen (no. 205) of this species in the New York state museum, taken on Grand Island, Niagara river, Aug. 1844. See 3d ann. rep't N. Y. state museum, p. 22; also Aub, 1857, 3:233-54.



- 101 (194) Great blue heron  
*Ardea herodias* Linn.

- 102 (196) American egret  
*Ardea egretta* Gmel.

- 103 (197) Snowy heron  
*Ardea candidissima* Gmel.

- 104 (199) Louisiana heron<sup>1</sup>  
*Ardea tricolor ruficollis* (Gosse)

---

<sup>1</sup> One Long Island record. See DeKay, 1843, p. 223. See also Giraud's *Birds of Long Island*, 1844, p. 281-82.



105 (200) Little blue heron  
*Ardea caerulea* Linn.

106 (201) Green heron  
*Ardea virescens* Linn.

107 (202) Black-crowned night heron  
*Nycticorax nycticorax naevius* (Bodd.)

108 (203) Yellow-crowned night heron  
*Nycticorax violaceus* (Linn.)



Order **Paludicolae**    *Cranes, rails, etc.*Family **GRUIDAE**    *Cranes*

- 109 (204) Whooping crane<sup>1</sup>  
    *Grus americana* (Linn.)

- 110 (206) Sandhill crane<sup>1</sup>  
    *Grus mexicana* (Müll.)

Family **RALLIDAE**    *Rails, gallinules and coots*

- 111 (208) King rail  
    *Rallus elegans* Aud.

- 112 (211) Clapper rail  
    *Rallus crepitans* Gmel.

---

<sup>1</sup> Historic in New York.





- 113 (212) Virginia rail  
Rallus virginianus Linn.
- 114 (214) Sora  
Porzana carolina (Linn.)
- 115 (215) Yellow rail  
Porzana noveboracensis (Gmel.)
- 116 (216) Black rail  
Porzana jamaicensis (Gmel.)



117 [217] Corn crake

*Crex crex* (Linn.)

118 (218) Purple gallinule

*Ionornis martinica* (Linn.)

119 (219) Florida gallinule

*Gallinula galeata* (Licht.)

120 (221) American coot

*Fulica americana* Gmel.



Order **Limicolae** *Shore birds*Family PHALAROPODIDAE *Phalaropes*

- 121 (222) Red phalarope  
*Crymophilus fulicarius* (Linn.)

- 122 (223) Northern phalarope  
*Phalaropus lobatus* (Linn.)

- 123 (224) Wilson's phalarope  
*Steganopus tricolor* Vieill.

Family RECURVIROSTRIDAE *Avocets and stilts*

- 124 (225) American avocet  
*Recurvirostra americana* Gmel.



- 125 (226) Black-necked stilt  
Himantopus mexicanus (Müll.)

Family SCOLOPACIDAE    *Snipes, sandpipers, etc.*

- 126 (228) American woodcock  
Philohela minor (Gmel.)

- 127 (230) Wilson's snipe  
Gallinago delicata (Ord)

- 128 (231) Dowitcher  
Macrorhamphus griseus (Gmel.)





- 129 (232) Long-billed dowitcher  
*Macrorhamphus scolopaceus* (Say)

- 130 (233) Stilt sandpiper  
*Micropalma himantopus* (Bonap.)

- 131 (234) Knot  
*Tringa canutus* Linn.

- 132 (235) Purple sandpiper  
*Tringa maritima* Brünn.



- 133 (239) Pectoral sandpiper  
*Tringa maculata* Vieill.
- 134 (240) White-rumped sandpiper  
*Tringa fuscicollis* Vieill.
- 135 (241) Baird's sandpiper  
*Tringa bairdii* (Coues)
- 136 (242) Least sandpiper  
*Tringa minutilla* Vieill.



- 137 [243] Dunlin<sup>1</sup>  
*Tringa alpina* Linn.

- 138 (243a) Red-backed sandpiper  
*Tringa alpina pacifica* (Coues)

- 139 (244) Curlew sandpiper<sup>2</sup>  
*Tringa ferruginea* Bünin.

- 140 (246) Semipalmated sandpiper  
*Ereunetes pusillus* (Linn.)

---

<sup>1</sup> One Long Island record. See *Auk*, 10: 78.

<sup>2</sup> See Coues's, *New England bird life*, 2:224; *Auk*, 1884, 1:32-33.



- 141 (247) Western sandpiper  
*Ereunetes occidentalis* Lawr.

- 142 (248) Sanderling  
*Calidris arenaria* (Linn.)

- 143 (249) Marbled godwit  
*Limosa fedoa* (Linn.)

- 144 (251) Hudsonian godwit  
*Limosa haemastica* (Linn.)





- 145 (254) Greater yellow-legs  
Totanus melanoleucus (Gmel.)

- 146 (255) Yellow-legs  
Totanus flavipes (Gmel.)

- 147 (256) Solitary sandpiper  
Helodromas solitarius (Wils.)

- 148 (258) Willet  
Symphemia semipalmata (Gmel.)



149 [260] Ruff<sup>1</sup>

*Pavoncella pugnax* (Linn.)

150 (261) Bartramian sandpiper

*Bartramia longicauda* (Bechst.)

151 (262) Buff-breasted sandpiper

*Tryngites subruficollis* (Vieill.)

152 (263) Spotted sandpiper

*Actitis macularia* (Linn.)

---

<sup>1</sup> Long Island records. See *Ann. lyc. nat. hist. N. Y.* 5: 220.



- 153 (264) Long-billed curlew  
*Numenius longirostris* Wils.
- 154 (265) Hudsonian curlew  
*Numenius hudsonicus* Lath.
- 155 (266) Eskimo curlew  
*Numenius borealis* (Forst.)
- 156 (H.L. 11.1) European curlew<sup>1</sup>  
*Numenius arquatus* (Linn.)

---

<sup>1</sup> There is in the New York state museum, (no. 324 in coll.) one individual of this species believed to have been taken on Long Island in 1853 and we think the evidence is sufficient to warrant its being included in the list of birds actually taken in New York. See *Auk*, 9: 390-92.



Family CHARADRIIDAE    *Plovers*157<sup>1</sup> [269] Lapwing<sup>1</sup>*Vanellus vanellus* (Linn.)

158 (270) Black-bellied plover

*Squatarola squatarola* (Linn.)

159 (272) American golden plover

*Charadrius dominicus* Müll.

160 (273) Killdeer

*Aegialitis vocifera* (Linn.)

---

<sup>1</sup> One specimen of this European form has been taken on Long Island. See *Auk*, 1886, 3: 438.





- 161 (274) Semipalmated plover  
*Aegialitis semipalmata* Bonap.
- 162 (277) Piping plover  
*Aegialitis meloda* (Ord)
- 163 (277a) Belted piping plover  
*Aegialitis meloda circumcincta* Ridgw.
- 164 (280) Wilson's plover  
*Aegialitis wilsonia* (Ord)



Family **APHRIZIDAE** *Surf birds and turnstones*

- 165 (283) Turnstone  
*Arenaria interpres* (Linn.)

Family **HAEMATOPODIDAE** *Oyster-catchers*

- 166 (286) American oyster-catcher  
*Haematopus palliatus* Temm.

Order **Gallinae** *Gallinaceous birds*Family **TETRAONIDAE** *Grouse, partridges, etc.*

- 167 (289) Bob-white  
*Colinus virginianus* (Linn.)

- 168 (298) Canada grouse  
*Canachites canadensis* (Linn.)



- 169 (300) Ruffed grouse  
Bonasa umbellus (Linn.)
- 170 (300a) Canadian ruffed grouse  
Bonasa umbellus togata (Linn.)
- 171 (301) Willow ptarmigan<sup>1</sup>  
Lagopus lagopus (Linn.)
- 172 (306) Heath hen<sup>2</sup>  
Tympanuchus cupido (Linn.)

---

<sup>1</sup> Watson, Lewis co. record. See *Bul. Nutt. orn. club*, 3: 41; 6: 233.

<sup>2</sup> Historic in New York state.



Family PHASIANIDAE *Pheasants, etc.*

- 173 (310a) Wild turkey<sup>1</sup>  
Meleagris gallopavo fera (Vieill.)

Order Columbæ *Pigeons*Family COLUMBIDAE *Pigeons*

- 174 (315) Passenger pigeon  
Ectopistes migratorius (Linn.)

- 175 (316) Mourning dove  
Zenaidura macroura (Linn.)

Order Raptores *Birds of prey*Family CATHARTIDAE *American vultures*

- 176 (325) Turkey vulture  
Cathartes aura (Linn.)

---

<sup>1</sup> Historic in New York state.





- 177 (326) Black vulture  
Catharista urubu (Vieill.)

Family FALCONIDAE    *Vultures, falcons, hawks, eagles, etc.*

- 178 (327) Swallow-tailed kite  
Elanoides forficatus (Linn.)

- 179 (331) Marsh hawk  
Circus hudsonius (Linn.)

- 180 (332) Sharp-shinned hawk  
Accipiter velox (Wils.)



- 181 (333) Cooper's hawk  
*Accipiter cooperii* (Bonap.)

- 182 (334) American goshawk  
*Accipiter atricapillus* (Wils.)

- 183 (337) Red-tailed hawk  
*Buteo borealis* (Gmel.)

- 184 (339) Red-shouldered hawk  
*Buteo lineatus* (Gmel.)



185 (342) Swainson's hawk<sup>1</sup>

*Buteo swainsoni* Bonap.

186 (343) Broad-winged hawk

*Buteo latissimus* (Wils.)

187 (347a) American rough-legged hawk

*Archibuteo lagopus sancti-johannis* (Gmel.)

188 (349) Golden eagle

*Aquila chrysaëtos* (Linn.)

---

<sup>1</sup> Cornwall N. Y. record. See *Auk*, 10: 83-84.



- 189 (352) Bald eagle  
*Haliaeetus leucocephalus* (Linn.)

- 190 (353) White gyrfalcon<sup>1</sup>  
*Falco islandus* Brünn.

- 191 (354a) Gyrfalcon<sup>2</sup>  
*Falco rusticolus gyrfalco* (Linn.)

- 192 (354b) Black gyrfalcon  
*Falco rusticolus obsoletus* (Gmel.)

---

<sup>1</sup> One taken on Long Island in 1856. See *Auk*, 10: 274-75; *Ann. l'yc. nat. hist. N. Y.* 8: 290.

<sup>2</sup> Long Island, fall of 1875.—*Lawrence*





193 (356) Duck hawk

*Falco peregrinus anatum* (Bonap.)

194 (357) Pigeon hawk

*Falco columbarius* Linn

195 (360) American sparrow hawk

*Falco sparverius* Linn.

196 (364) American osprey

*Pandion haliaëtus carolinensis* (Gmel.)



Family STRIGIDAE    *Barn owls*

- 197 (365) American barn owl  
    *Strix pratincta* Bonap.

Family BUBONIDAE    *Horned owls, etc.*

- 198 (366) American long-eared owl  
    *Asio wilsonianus* (Less.)

- 199 (367) Short-eared owl  
    *Asio accipitrinus* (Pall.)

- 200 (368) Barred owl  
    *Syrnium nebulosum* (Forst.)



201 (370) Great gray owl  
    *Scotiaptex cinerea* (Gmel.)

202 (372) Saw-whet owl  
    *Nyctala acadica* (Gmel.)

203 (373) Screech owl  
    *Megascops asio* (Linn.)

204 (375) Great horned owl  
    *Bubo virginianus* (Gmel.)



- 205 (376) Snowy owl  
Nyctea nyctea (Linn.)

- 206 (377a) American hawk owl  
Surnia ulula caparoch (Müll.)

- 207 (378) Burrowing owl<sup>1</sup>  
Speotyto cunicularia hypogaea (Bonap.)

Order **Psittaci**    *Parrots, etc.*

Family **PSITTACIDAE**    *Parrots*

- 208 (382) Carolina paroquet<sup>2</sup>  
Conurus carolinensis (Linn.)

---

<sup>1</sup> One New York record. See *Forest and stream*, 12 Aug. 1875, 5: 4.

<sup>2</sup> Historic in New York state.





Order **Coccyges** *Cuckoos, etc.*Family CUCULIDAE *Cuckoos, etc.*

- 209 (387) Yellow-billed cuckoo  
Coccyzus americanus (Linn.)

- 210 (388) Black billed cuckoo  
Coccyzus erythrophthalmus (Wils.)

Family ALCEDINIDAE *Kingfishers*

- 211 (390) Belted kingfisher  
Ceryle alcyon (Linn.)

Order **Pici** *Woodpeckers, etc.*Family PICIDAE *Woodpeckers*

- 212 (393) Hairy woodpecker  
Dryobates villosus (Linn.)



- 213 (394c) Downy woodpecker  
*Dryobates pubescens medianus* (Swains.)

- 214 (400) Arctic three-toed woodpecker  
*Picoides arcticus* (Swains.)

- 215 (401) American three-toed woodpecker  
*Picoides americanus* Brehm

- 216 (402) Yellow-bellied sapsucker  
*Sphyrapicus varius* (Linn.)



- 217 (405a) Northern pileated woodpecker  
*Ceophloeus pileatus abieticola* Bangs

- 218 (406) Red-headed woodpecker  
*Melanerpes erythrocephalus* (Linn.)

- 219 (409) Red-bellied woodpecker  
*Melanerpes carolinus* (Linn.)

- 220 (412a) Northern flicker  
*Colaptes auratus luteus* Bangs



Order **Machrochires**    *Goatsuckers, swifts, etc.*Family CAPRIMULGIDAE    *Goatsuckers, etc.*

- 221 (417) Whip-poor-will  
    *Antrostomus vociferus* (Wils.)

- 222 (420) Nighthawk  
    *Chordeiles virginianus* (Gmel.)

Family MICROPODIDAE    *Swifts*

- 223 (423) Chimney swift  
    *Chaetura pelagica* (Linn.)

Family TROCHILIDAE    *Hummingbirds*

- 224 (428) Ruby-throated hummingbird  
    *Trochilus colubris* Linn.





Order **Passeres** *Perching birds*Family **TYRANNIDAE** *Tyrant flycatchers*

- 225 (444) Kingbird  
Tyrannus tyrannus (Linn.)

- 226 (447) Arkansas kingbird  
Tyrannus verticalis Say

- 227 (452) Crested flycatcher  
Myiarchus crinitus (Linn.)

- 228 (456) Phoebe  
Sayornis phoebe (Lath.)



229 (459) Olive-sided flycatcher  
*Contopus borealis* (Swains.)

230 (461) Wood pewee  
*Contopus virens* (Linn.)

231 (463) Yellow-bellied flycatcher  
*Empidonax flaviventris* Baird

232 (465) Green-crested flycatcher  
*Empidonax virescens* (Vieill.)



- 233 (466a) Alder flycatcher  
*Empidonax traillii alnorum* Brewst.

- 234 (467) Least flycatcher  
*Empidonax minimus* Baird

Family ALAUDIDAE    *Larks*

- 235 (474) Horned lark  
*Otocoris alpestris* (Linn.)
- 236 (474b) Prairie horned lark  
*Otocoris alpestris praticola* Hensh.



Family CORVIDAE    *Crows, jays, magpies, etc.*

237 (477) Blue jay

*Cyanocitta cristata* (Linn.)

238 (484) Canada jay

*Perisoreus canadensis* (Linn.)

239 (486a) Northern raven

*Corvus corax principalis* Ridgw.

240 (488) American crow

*Corvus americanus* Aud.

1. *Amphispiza bilineata* (Aud.)

2. *Amphispiza bilineata* (Aud.)

3. *Amphispiza bilineata* (Aud.)

4. *Amphispiza bilineata* (Aud.)

5. *Amphispiza bilineata* (Aud.)



- 241 (490) Fish crow  
Corvus ossifragus Wils.

Family ICTERIDAE    *Blackbirds, orioles, etc.*

- 242 (494) Bobolink  
Dolichonyx oryzivorus (Linn.)

- 243 (495) Cowbird  
Molothrus ater (Bodd.)

- 244 (498) Red-winged blackbird  
Agelaius phoeniceus (Linn.)



245 (501) Meadowlark

*Sturnella magna* (Linn.)

246 (506) Orchard oriole

*Icterus spurius* (Linn.)

247 (507) Baltimore oriole

*Icterus galbula* (Linn.)

248 (509) Rusty blackbird

*Scolecophagus carolinus* (Müll.)



- 249 (511) Purple grackle  
Quiscalus quiscula (Linn.)

- 250 (511b) Bronzed grackle  
Quiscalus quiscula aeneus (Ridgw.)

Family FRINGILLIDAE *Finches, sparrows, etc.*

- 251 (514) Evening grosbeak  
Coccothraustes vespertinus (Coop.)

- 252 (515) Pine grosbeak  
Pinicola enucleator canadensis (Cab.)



- 253 (517) Purple finch  
Carpodacus purpureus (Gmel.)
- 254 (521) American crossbill  
Loxia curvirostra minor (Brehm)
- 255 (522) White-winged crossbill  
Loxia leucoptera Gmel.
- 256 (528) Redpoll  
Acanthis linaria (Linn.)





- 257 (528b) Greater redpoll  
*Acanthis linaria rostrata* (Coues)

- 258 (529) American goldfinch  
*Astragalinus tristis* (Linn.)

- 259 (533) Pine siskin  
*Spinus pinus* (Wils.)

- 260 (534) Snowflake  
*Passerina nivalis* (Linn.)



261 (536) Lapland longspur  
*Calcarius lapponicus* (Linn.)

262 (538) Chestnut-collared longspur<sup>1</sup>  
*Calcarius ornatus* (Townsend)

263 (540) Vesper sparrow  
*Poecetes gramineus* (Gmel.)

264 (541) Ipswich sparrow  
*Ammodramus princeps* (Mayn.)

---

<sup>1</sup> Long Island record. See *Auk*, 6: 190.



- 265 (542a) Savanna sparrow  
*Ammodramus sandwichensis savanna* (Wils.)

- 266 (546) Grasshopper sparrow  
*Ammodramus savannarum passerinus* (Wils.)

- 267 (547) Henslow's sparrow  
*Ammodramus henslowii* (Aud.)

- 268 (548) Leconte's sparrow<sup>1</sup>  
*Ammodramus leconteii* (Aud.)

---

<sup>1</sup> This is included on the authority of Mr Louis Agassiz Fuertes, who reports the capture of a young bird of this species at Ithaca, 11 Oct. 1897. See *Auk*, 1898, 15: 188-89.



- 269 (549) Sharp-tailed sparrow  
*Ammodramus caudacutus* (Gmel.)
- 270 (549.1) Nelson's sparrow  
*Ammodramus nelsoni* (Allen)
- 271 (549.1a) Acadian sharp-tailed sparrow  
*Ammodramus nelsoni subvirgatus* (Dwight)
- 272 (550) Seaside sparrow  
*Ammodramus maritimus* (Wils.)





273 (552) Lark sparrow

*Chondestes grammacus* (Say)

274 (554) White-crowned sparrow

*Zonotrichia leucophrys* (Forst.)

275 (558) White-throated sparrow

*Zonotrichia albicollis* (Gmel.)

276 (559) Tree sparrow

*Spizella monticola* (Gmel.)



277 (560) Chipping sparrow  
*Spizella socialis* (Wils.)

278 (563) Field sparrow  
*Spizella pusilla* (Wils.)

279 (567) Slate-colored junco  
*Junco hyemalis* (Linn.)

280 (581) Song sparrow  
*Melospiza fasciata* (Gmel.)<sup>1</sup>

---

<sup>1</sup> Oberholser has recently shown that the species name *fasciata* is untenable for this form and that *melodia* should be adopted. See *Auk*, 16: 182-83.



- 281 (583) Lincoln's sparrow  
    *Melospiza lincolnii* (Aud.)
- 282 (584) Swamp sparrow  
    *Melospiza georgiana* (Lath.)
- 283 (585) Fox sparrow  
    *Passerella iliaca* (Merr.)
- 284 (587) Towhee  
    *Pipilo erythrophthalmus* (Linn.)



285 (593) Cardinal  
*Cardinalis cardinalis* (Linn.)

286 (595) Rose-breasted grosbeak  
*Zamelodia ludoviciana* (Linn.)

287 (597) Blue grosbeak<sup>1</sup>  
*Guiraca caerulea* (Linn.)

288 (598) Indigo bunting  
*Cyanospiza cyanea* (Linn.)

---

<sup>1</sup> See De Kay, 1842, p. 146; *Bul. Nutt. orn. club*, 3:132; *Auk*, 10:276.





- 289 (601) Painted bunting<sup>1</sup>  
*Cyanospiza ciris* (Linn.)

- 290 (604) Dickcissel  
*Spiza americana* (Gmel.)

- 291 (605) Lark bunting<sup>2</sup>  
*Calamospiza melanocorys* Stejn.

Family TANAGRIDAE    *Tanagers*

- 292 (607) Louisiana tanager<sup>3</sup>  
*Piranga ludoviciana* (Wils.)

---

<sup>1</sup> See *Bul. Nutt. orn. club*, 3: 132; *Forest and stream*, 1884, p. 424.

<sup>2</sup> This western species has once been taken on Long Island. See *Auk*, 6: 192, 1889.

<sup>3</sup> One New York record. See *Auk*, 1890, 7: 55.



- 293 (608) Scarlet tanager  
Piranga erythromelas Vieill.

- 294 (610) Summer tanager  
Piranga rubra (Linn.)

Family HIRUNDINIDAE    *Swallows*

- 295 (611) Purple martin  
Progne subis (Linn.)

- 296 (612) Cliff swallow  
Petrochelidon lunifrons (Say)



- 297 (613) Barn swallow  
Hirundo erythrogaster Bodd.
- 298 (614) Tree swallow  
Tachycineta bicolor (Vieill.)
- 299 (616) Bank swallow  
Clivicola riparia (Linn.)
- 300 (617) Rough-winged swallow  
Stelgidopteryx serripennis (Aud.)



Family AMPELIDAE    *Waxwings, etc.*

- 301 (618) Bohemian waxwing  
    *Ampelis garrulus* Linn.

- 302 (619) Cedar waxwing  
    *Ampelis cedrorum* (Vieill.)

Family LANIIDAE<sup>1</sup>    *Shrikes*

- 303 (621) Northern shrike  
    *Lanius borealis* Vieill.

- 304 (622) Loggerhead shrike  
    *Lanius ludovicianus* Linn.

---

<sup>1</sup> Three species of shrikes have been recognized in New York state, but the status of the forms is not well understood, it often being a matter of difficulty to assign individual specimens to their proper species. William Palmer, after an examination of 176 specimens, proposes to "relegate *excubitorides* to the plains region west of the immediate Mississippi wooded drainage area, *ludovicianus* to the south Atlantic and gulf coasts, and Florida, and to recognize a new form (*L. l. migrans*) as occupying much of the remaining region of the east." The acceptance of his views has been deferred by the American ornithologists union for lack of material, and so it seems best for the present to let the species stand as given in *Auk*, 15: 241-58; also 16: 133.





- 305 (622a) White-rumped shrike  
*Lanius ludovicianus excubitorides* (Swains.)

Family VIREONIDAE    *Vireos*

- 306 (624) Red-eyed vireo  
*Vireo olivaceus* (Linn.)
- 307 (626) Philadelphia vireo  
*Vireo philadelphicus* (Cass.)
- 308 (627) Warbling vireo  
*Vireo gilvus* (Vieill.)



- 309 (628) Yellow-throated vireo  
Vireo flavifrons Vieill.

- 310 (629) Blue-headed vireo  
Vireo solitarius (Wils.)

- 311 (629b) Plumbeous vireo<sup>1</sup>  
Vireo solitarius plumbeus (Coues)

- 312 (631) White-eyed vireo  
Vireo noveboracensis (Gmel.)

---

<sup>1</sup> One New York record. See *Auk* 1894, 11:59.



Family MNIOTILTIDAE    *Wood warblers*

- 313 (636) Black and white warbler  
Mniotilta varia (Linn.)

- 314 (637) Prothonotary warbler  
Protonotaria citrea (Bodd.)

- 315 (639) Worm-eating warbler  
Helmitherus vermivorus (Gmel.)

- 316 (641) Blue-winged warbler  
Helminthophila pinus (Linn.)



- 317 (642) Golden-winged warbler  
*Helminthophila chrysoptera* (Linn.)

- 318 (H.L.20) Lawrence's warbler<sup>1</sup>  
*Helminthophila lawrencei* (Herrick)

- 319 (H.L.21) Brewster's warbler<sup>1</sup>  
*Helminthophila leucobronchialis* (Brewst.)

- 320 (645) Nashville warbler  
*Helminthophila rubricapilla* (Wils.)

---

<sup>1</sup> These forms are included in the hypothetical list of the American ornithologists union check list, as they are supposed to be hybrids between *H. pinus* and *H. chrysoptera*. Both have been taken in New York, and are here included to learn more of their abundance and status in our state. See *Auk*, 1883, 5: 427; *Bul. Nutt. orn. club*, 4: 234, 1879; *Auk*, 1892, 9: 306.





- 321 (646) Orange-crowned warbler  
*Helminthophila celata* (Say)
- 322 (647) Tennessee warbler  
*Helminthophila peregrina* (Wils.)
- 323 (648a) Northern parula warbler  
*Compsothlypis americana usneae* Brewst.
- 324 (650) Cape May warbler  
*Dendroica tigrina* (Gmel.)



- 325 (652) Yellow warbler  
*Dendroica aestiva* (Gmel.)
- 326 (654) Black-throated blue warbler  
*Dendroica caerulescens* (Gmel.)
- 327 (655) Myrtle warbler  
*Dendroica coronata* (Linn.)
- 328 (657) Magnolia warbler  
*Dendroica maculosa* (Gmel.)



- 329 (658) Cerulean warbler  
*Dendroica rara* Wils.

- 330 (659) Chestnut-sided warbler  
*Dendroica pensylvanica* (Linn.)

- 331 (660) Bay-breasted warbler  
*Dendroica castanea* (Wils.)

- 332 (661) Black-poll warbler  
*Dendroica striata* (Forst.)



- 333 (662) Blackburnian warbler  
*Dendroica blackburniae* (Gmel.)

- 334 (663) Yellow-throated warbler  
*Dendroica dominica* (Linn.)

- 335 (667) Black-throated green warbler  
*Dendroica virens* (Gmel.)

- 336 (671) Pine warbler  
*Dendroica vigorsii* (Aud.)





- 337 (672) Palm warbler  
Dendroica palmarum (Gmel.)
- 338 (672a) Yellow palm warbler  
Dendroica palmarum hypochrysea Ridgw.
- 339 (673) Prairie warbler  
Dendroica discolor (Vieill.)
- 340 (674) Oven-bird  
Seiurus aurocapillus (Linn.)



- 341 (675) Water-thrush  
*Seiurus noveboracensis* (Gmel.)

- 342 (676) Louisiana water-thrush  
*Seiurus motacilla* (Vieill.)

- 343 (677) Kentucky warbler  
*Geothlypis formosa* (Wils.)

- 344 (678) Connecticut warbler  
*Geothlypis agilis* (Wils.)



345 (679) Mourning warbler  
*Geothlypis philadelphia* (Wils.)

346 (681) Maryland yellow-throat  
*Geothlypis trichas* (Linn.)

347 (683) Yellow-breasted chat  
*Icteria virens* (Linn.)

348 (684) Hooded warbler  
*Wilsonia mitrata* (Gmel.)



- 349 (685) Wilson's warbler  
    *Wilsonia pusilla* (Wils.)

- 350 (686) Canadian warbler  
    *Wilsonia canadensis* (Linn.)

- 351 (687) American redstart  
    *Setophaga ruticilla* (Linn.)

Family MOTACILLIDAE   *Wagtails*

- 352 (697) American pipit  
    *Anthus pensilvanicus* (Lath.)





Family TROGLODYTIDAE *Wrens, thrashers, etc.*

## 353 (703) Mockingbird

*Mimus polyglottos* (Linn.)

## 354 (704) Catbird

*Galeoscoptes carolinensis* (Linn.)

## 355 (705) Brown thrasher

*Harporhynchus rufus* (Linn.)

## 356 (718) Carolina wren

*Thryothorus ludovicianus* (Lath.)



- 357 (721) House wren  
*Troglodytes aëdon* Vieill.
- 358 (722) Winter wren  
*Anorthura hiemalis* (Vieill.)
- 359 (724) Short-billed marsh wren  
*Cistothorus stellaris* (Licht.)
- 360 (725) Long-billed marsh wren  
*Cistothorus palustris* (Wils.)



Family CERTHIIDAE *Creepers*

- 361 (726) Brown creeper  
*Certhia familiaris fusca* (Barton)<sup>1</sup>

Family PARIDAE *Nuthatches and tits*

- 362 (727) White-breasted nuthatch  
*Sitta carolinensis* Lath.

- 363 (728) Red-breasted nuthatch  
*Sitta canadensis* Linn.

- 364 (729) Brown-headed nuthatch<sup>1</sup>  
*Sitta pusilla* Lath.

---

<sup>1</sup> Oberholser has recently shown that this name is untenable and that the former *c. f. americana* (Bonap.) must be restored. See *Auk*, 16: 185.

<sup>2</sup> One New York record. See *Auk*, 1888, 5:432.



- 365 (731) Tufted titmouse  
Parus bicolor Linn.

- 366 (735) Chickadee  
Parus atricapillus Linn.

- 367 (740) Hudsonian chickadee  
Parus hudsonicus Forst.

Family SYLVIIDAE *Warblers, kinglets, gnatcatchers*

- 368 (748) Golden-crowned kinglet  
Regulus satrapa Licht.





- 369 (749) Ruby-crowned kinglet  
Regulus calendula (Linn.)

- 370 (751) Blue-gray gnatcatcher  
Polioptila caerulea (Linn.)

Family TURDIDAE *Thrushes, stone-chats, blue-birds, etc.*

- 371 (755) Wood thrush  
Hylocichla mustelina (Gmel.)

- 372 (756) Wilson's thrush  
Hylocichla fuscescens (Steph.)



- 373 (757) Gray-cheeked thrush  
*Hylocichla aliciae* (Baird)

- 374 (757a) Bicknell's thrush  
*Hylocichla aliciae bicknelli* Ridgw.

- 375 (758a) Olive-backed thrush  
*Hylocichla ustulata swainsonii* (Cab.)

- 376 (759b) Hermit thrush  
*Hylocichla aonalaschkae pallasii* (Cab.)



- 377 (761) American robin  
    *Merula migratoria* (Linn.)
- 378 (763) Varied thrush<sup>1</sup>  
    *Hesperocichla naevia* (Gmel.)
- 379 (765) Wheatear<sup>2</sup>  
    *Saxicola oenanthe* (Linn.)
- 380 (766) Bluebird  
    *Sialia sialis* (Linn.)

---

<sup>1</sup> "There are no less than three valid records for this species near New York city." See Dwight. *Auk*, 12:70; *Proc. Boston soc. nat. hist.* 1848, 3:17; *Ann. lyc. nat. hist. N. Y.* 1867, 8:281; Coues's *Birds Colorado valley* 1878, p. 19; Baird Brewer and Ridgway's *Birds*, 1874, 1:29.

<sup>2</sup> Long Island records. See *Auk*, 3:490; also 10:277; *Ann. lyc. nat. hist. N. Y.* 8: 282, 1867.



INTRODUCED SPECIES<sup>1</sup>Family ALAUDIDAE *Larks*381 [473] Skylark<sup>2</sup>*Alauda arvensis* Linn.Family STURNIDAE *Starlings*382 [493] Starling<sup>3</sup>*Sturnus vulgaris* Linn.Family FRINGILLIDAE *Finches, sparrows, etc.*

383 European goldfinch

*Carduelis carduelis* (Linn.)

384 House sparrow

*Passer domesticus* (Linn.)

---

<sup>1</sup> Exclusive of game birds.

<sup>2</sup> This European form is accidental in Greenland and Bermuda. Has been introduced in this country and in 1887 was found breeding at Flatbush L. I. See *Auk*, 5: 180; 12:390.

<sup>3</sup> See *Auk*, 15: 55-56.





## PART 2

## HYPOTHETIC LIST

Including those forms, of which we have no satisfactory record, but which, having been taken in adjoining states, may be looked for in New York.

---

- 1 (28) Mandt's guillemot  
Cepphus mandtii (Licht.)

Arctic regions . . . south . . . in winter, to Mass.—*American ornithologists union*

- 2 (30) Murre  
Uria troile (Linn.)

Southward on the coast of North America, in winter, to southern New England.—*American ornithologists union*. This species has been reported from our state, but it seems probable that the specimens taken were immature individuals of *U. lumme*.

- 3 (67) Cabot's tern  
Sterna sandvicensis acufflvida (Cabot.)

South Atlantic and gulf coasts of the United States, north, casually, to southern New England.—*American ornithologists union*

- 4 (86) Fulmar  
Fulmarus glacialis (Linn.)

North Atlantic, south on the American side to the coast of Massachusetts, casually to New Jersey.—*American ornithologists union*. Accidental in northern New Jersey.—*Chapman*



## 5 (86a) Lesser fulmar

Fulmarus glacialis minor (Kjaerboelling)

North Atlantic, south on American side to coast of Massachusetts.—*American ornithologists union*

## 6 (126) Brown pelican

Pelecanus fuscus (Linn.)

Occasionally strays to Illinois and Massachusetts.—Chapman, *Birds of eastern North America*. De Kay (1842), mentions this form as "an occasional visitor to the coast of our state," but gives no records.—*Birds*, p. 294

## 7 [138] European teal

Nettion crecca (Linn.)

Occasional in eastern North America.—*American ornithologists union*. New Jersey, Connecticut, and Massachusetts records.—Abbott, *Geology of New Jersey*, p. 792; Treat, *Auk*, 8: 112; Mackay, *Auk*, 7: 294

## 8 (141) Cinnamon teal

Querquedula cyanoptera (Vieill.)

Accidental east of the Mississippi.



## 9 (173a) White-bellied brant

*Branta bernicla glaucogastra* (Brehm.)

This subspecies has been recognized in America by Coues. See *Auk*, 14 : 207-8.  
This form may be looked for in our state, though I can find no records as yet.

## 10 (191.1) Cory's least bittern

*Ardetta neoxena* Cory

Michigan and Ontario records. Will undoubtedly be found in our state.

## 11 [227] European woodcock

*Scolopax rusticola* (Linn.)

Accidental in the United States. There are New Jersey, Pennsylvania and Rhode Island records; and we may expect to find this form in New York.

## 12 (371) Richardson's owl

*Nyctala tengmalmi richardsoni* (Bonap.)

South in winter into the northern United States. *American ornithologist union*. Rhode Island record. See *Auk*, 3 : 271



- 13 (395) Red-cockaded woodpecker  
*Dryobates borealis* (Vieill.)

"A southern United States form has been taken as far north as Hoboken, New Jersey." See Lawrence, *Ann. lyc. nat. hist. N. Y.*, 8: 291.

- 14 (443) Scissor-tailed flycatcher  
*Milvulus forficatus* (Gmel.)

A southwestern United States form that has occurred accidentally in New Jersey and New England.

- 15 (475) American magpie.  
*Pica pica hudsonica* (Lab.)

De Kay (1842) mentions it as having occurred in the northwestern part of the state near Niagara, but does not give any records.

- 16 (497) Yellow-headed blackbird  
*Xanthocephalus xanthocephalus* (Bonap.)

Accidental in Ontario and the Atlantic states, Massachusetts, etc.—*American ornithologists union*





- 17 (736) Carolina chickadee  
Parus carolinensis Aud.

Princeton N. J., is usually considered its northern limit. See Scott's *The country*, 1878, 1: 334.



# INDEX

The superior figures tell the exact place on the page in ninths; e. g. 201<sup>8</sup> means page 201 beginning in the third ninth of the page, i. e. about one third of the way down.

- Acanthis* linaria, 323<sup>7</sup>  
     rostrata, 325<sup>1</sup>  
*Accipiter* atricapillus, 287<sup>8</sup>  
     cooperii, 287<sup>1</sup>  
     velox, 285<sup>7</sup>  
*Actitis* macularia, 271<sup>7</sup>  
*Aegialitis* meloda, 277<sup>8</sup>  
     circumcincta, 277<sup>5</sup>  
     semipalmata, 277<sup>1</sup>  
     vocifera, 275<sup>7</sup>  
     wilsonia, 277<sup>7</sup>  
*Aestrelata* hasitata, 217<sup>7</sup>  
     scalaris, 219<sup>1</sup>  
*Agelaius* phoeniceus, 317<sup>7</sup>  
*Aix* sponsa, 229<sup>5</sup>  
*Alauda* arvensis, 387<sup>1</sup>  
*Alaudidae*, 313<sup>5</sup>, 387<sup>1</sup>  
*Alca* torda, 201<sup>3</sup>  
*Alcedinidae*, 301<sup>5</sup>  
*Alcidae*, 199<sup>5</sup>-201<sup>5</sup>  
*Alle alle*, 201<sup>5</sup>  
*Ammodramus* caudacutus, 331<sup>1</sup>  
     henslowii, 329<sup>5</sup>  
     leconteii, 329<sup>7</sup>  
     maritimus, 331<sup>7</sup>  
     nelsoni, 331<sup>8</sup>  
     subvirgatus, 331<sup>5</sup>  
     princeps, 327<sup>7</sup>  
     sandwichensis savanna, 329<sup>1</sup>  
     savannarum passerinus, 329<sup>8</sup>  
*Ampelidae*, 347<sup>1</sup>  
*Ampelis* cedrorum, 347<sup>8</sup>  
     garrulus, 347<sup>1</sup>  
*Anas* boschas, 225<sup>3</sup>  
     obscura, 225<sup>5</sup>  
*Anatidae*; 223<sup>5</sup>-43<sup>5</sup>  
*Anorthura* hiemalis, 375<sup>7</sup>  
*Anser* albifrons gambeli, 239<sup>7</sup>  
*Anseres*, 223<sup>5</sup>-43<sup>5</sup>  
*Anthus* pensilvanicus, 371<sup>7</sup>  
*Antrostomus* vociferus, 307<sup>7</sup>  
*Aphrizidae*, 279<sup>1</sup>  
*Aquila* chrysaetos, 289<sup>7</sup>  
*Archibuteo* lagopus sancti-johannis,  
     289<sup>5</sup>  
*Ardea* caerulea, 249<sup>1</sup>  
     candidissima, 247<sup>5</sup>  
     egretta, 247<sup>8</sup>  
     herodias, 247<sup>1</sup>  
     tricolor ruficollis, 247<sup>7</sup>  
     virescens, 249<sup>3</sup>  
*Ardeidae*, 245<sup>5</sup>-49<sup>9</sup>  
*Ardetta* exilis, 245<sup>7</sup>  
     neoxena, 393<sup>3</sup>  
*Arenaria* interpres, 279<sup>1</sup>  
*Asio* accipitrinus, 295<sup>5</sup>  
     wilsonianus, 295<sup>3</sup>  
*Astragalinus* tristis, 325<sup>5</sup>  
*Auk*, razor-billed, 201<sup>8</sup>  
*Avocet*, American, 257<sup>8</sup>  
*Aythya* affinis, 231<sup>7</sup>  
     americana, 231<sup>1</sup>  
     collaris, 233<sup>1</sup>  
     marila, 231<sup>5</sup>  
     vallisneria, 231<sup>3</sup>

- Baldpate**, 227<sup>3</sup>  
*Bartramia longicauda*, 271<sup>3</sup>  
 Birds of prey, 283<sup>7</sup>-99<sup>6</sup>  
*Bittern*, American, 245<sup>5</sup>  
     Cory's least, 393<sup>3</sup>  
     least, 245<sup>7</sup>  
*Blackbird*, red-winged, 317<sup>7</sup>  
     rusty, 319<sup>7</sup>  
     yellow-headed, 395<sup>7</sup>  
*Bluebird*, 385<sup>7</sup>  
*Bobolink*, 317<sup>3</sup>  
*Bob-white*, 279<sup>6</sup>  
*Bonasa umbellus*, 281<sup>1</sup>  
     *togata*, 281<sup>3</sup>  
*Booby*, 221<sup>1</sup>  
*Botaurus lentiginosus*, 245<sup>5</sup>  
*Brant*, 241<sup>5</sup>  
     black, 241<sup>7</sup>  
     white-bellied, 393<sup>1</sup>  
*Branta bernicla*, 241<sup>5</sup>  
     *glaucogastra*, 393<sup>1</sup>  
     *canadensis*, 241<sup>1</sup>  
     *leucopsis*, 243<sup>1</sup>  
         *hutchinsii*, 241<sup>3</sup>  
         *nigracans*, 241<sup>7</sup>  
*Bubo virginianus*, 297<sup>7</sup>  
*Bubonidae*, 295<sup>3</sup>-99<sup>6</sup>  
*Buffle-head*, 233<sup>7</sup>  
*Bunting*, indigo, 339<sup>7</sup>  
     lark, 341<sup>5</sup>  
     painted, 341<sup>1</sup>  
*Buteo borealis*, 287<sup>5</sup>  
     *latissimus*, 289<sup>3</sup>  
     *lineatus*, 287<sup>7</sup>  
     *swainsoni*, 289<sup>1</sup>  
**Calamospiza melanocorys**, 341<sup>5</sup>  
*Calcarius lapponicus*, 327<sup>1</sup>  
     *ornatus*, 327<sup>3</sup>  
*Calidris arenaria*, 267<sup>3</sup>  
*Camptolaimus labradorius*, 235<sup>5</sup>  
*Canachites canadensis*, 279<sup>7</sup>  
*Canvas-back*, 231<sup>3</sup>  
*Caprimulgidae*, 307<sup>1</sup>  
*Cardinal*, 339<sup>1</sup>  
*Cardinalis cardinalis*, 339<sup>1</sup>  
*Carduelis carduelis*, 387<sup>5</sup>  
*Carpodacus purpureus*, 323<sup>1</sup>  
*Catbird*, 373<sup>7</sup>  
*Catharista urubu*, 285<sup>1</sup>  
*Cathartes aura*, 283<sup>7</sup>  
*Cathartidae*, 283<sup>7</sup>-85<sup>3</sup>  
*Ceophloeus pileatus abieticola*, 305<sup>1</sup>  
*Cephus grylle*, 199<sup>7</sup>  
     *mandtii*, 389<sup>2</sup>  
*Certhia familiaris fusca*, 377<sup>1</sup>  
*Certhiidae*, 377<sup>1</sup>  
*Ceryle alcyon*, 301<sup>5</sup>  
*Chaetura pelagica*, 307<sup>5</sup>  
*Charadriidae*, 275<sup>1</sup>-77<sup>9</sup>  
*Charadrius dominicus*, 275<sup>5</sup>  
*Charitonetta albeola*, 233<sup>7</sup>  
*Chat*, yellow-breasted, 369<sup>6</sup>  
*Chauleasmus streperus*, 225<sup>7</sup>  
*Chen caerulescens*, 239<sup>5</sup>  
     *hyperborea nivalis*, 239<sup>3</sup>  
*Chickadee*, 379<sup>3</sup>  
     *Carolina*, 397<sup>1</sup>  
     *Hudsonian*, 379<sup>5</sup>  
*Chondestes grammacus*, 333<sup>1</sup>  
*Chordeiles virginianus*, 307<sup>3</sup>  
*Ciconiidae*, 245<sup>3</sup>  
*Circus hudsonius*, 285<sup>5</sup>  
*Cistothorus palustris*, 375<sup>7</sup>  
     *stellaris*, 375<sup>5</sup>  
*Clangula clangula americana*, 233<sup>3</sup>  
     *islandica*, 233<sup>5</sup>  
*Clivicola riparia*, 345<sup>5</sup>  
*Coccythraustes vespertinus*, 321<sup>5</sup>  
*Coccyges*, 301<sup>1</sup>  
*Coccyzus americanus*, 301<sup>2</sup>  
     *erythrophthalmus*, 301<sup>3</sup>  
*Colaptes auratus luteus*, 305<sup>7</sup>  
*Colinus virginianus*, 279<sup>6</sup>  
*Columbae*, 283<sup>3</sup>  
*Columbidae*, 283<sup>3</sup>  
*Colymbus auritus*, 197<sup>4</sup>  
     *holboellii*, 197<sup>3</sup>  
*Compsothlypis americana usneae*,  
     357<sup>5</sup>  
*Contopus borealis*, 311<sup>1</sup>  
     *virens*, 311<sup>3</sup>

- Conurus carolinensis*, 299<sup>7</sup>  
 Coot, American, 255<sup>7</sup>  
 Cormorant, 221<sup>5</sup>  
     double-crested, 221<sup>7</sup>  
 Corvidae, 315<sup>1-17</sup><sup>3</sup>  
*Corvus americanus*, 315<sup>7</sup>  
     *corax principalis*, 315<sup>5</sup>  
     *ossifragus*, 317<sup>1</sup>  
 Cowbird, 317<sup>5</sup>  
 Crane, corn, 255<sup>1</sup>  
 Crane, sandhill, 251<sup>7</sup>  
     whooping, 251<sup>2</sup>  
 Creeper, brown, 377<sup>1</sup>  
*Crex crex*, 255<sup>1</sup>  
 Crossbill, American, 323<sup>3</sup>  
     white-winged, 323<sup>5</sup>  
 Crow, American, 315<sup>7</sup>  
     fish, 317<sup>1</sup>  
*Crymophilus fulicarius*, 257<sup>1</sup>  
 Cuckoo, black billed, 301<sup>3</sup>  
     yellow-billed, 301<sup>2</sup>  
 Cuculidae, 301<sup>1</sup>  
 Curlew, Eskimo, 273<sup>5</sup>  
     European, 273<sup>7</sup>  
     Hudsonian, 273<sup>3</sup>  
     long-billed, 273<sup>1</sup>  
*Cyanocitta cristata*, 315<sup>1</sup>  
*Cyanospiza ciris*, 341<sup>1</sup>  
     *cyanea*, 339<sup>7</sup>  
**Dafila** *acuta*, 229<sup>3</sup>  
*Dendroica aestiva*, 359<sup>1</sup>  
     *blackburniae*, 363<sup>1</sup>  
     *caerulescens*, 359<sup>3</sup>  
     *castanea*, 361<sup>5</sup>  
     *coronata*, 359<sup>5</sup>  
     *discolor*, 365<sup>5</sup>  
     *dominica*, 363<sup>3</sup>  
     *maculosa*, 359<sup>7</sup>  
     *palmarum*, 365<sup>1</sup>  
         *hypochrysea*, 365<sup>3</sup>  
     *pensylvanica*, 361<sup>3</sup>  
     *rara*, 361<sup>1</sup>  
     *striata*, 361<sup>7</sup>  
     *tigrina*, 357<sup>7</sup>  
     *vigorsii*, 363<sup>7</sup>  
         *viens*, 363<sup>5</sup>  
 Dickcissel, 341<sup>3</sup>  
 Diving birds, 197<sup>2-201</sup><sup>6</sup>  
*Dolichonyx oryzivorus*, 317<sup>3</sup>  
 Dove, mourning, 283<sup>5</sup>  
 Dovekie, 201<sup>5</sup>  
 Dowitcher, 259<sup>7</sup>  
     long-billed, 261<sup>1</sup>  
*Dryobates borealis*, 395<sup>1</sup>  
     *pubescens medianus*, 303<sup>1</sup>  
     *villosus*, 301<sup>3</sup>  
 Duck, American scaup, 231<sup>5</sup>  
     black, 225<sup>5</sup>  
     harlequin, 235<sup>3</sup>  
     Labrador, 235<sup>5</sup>  
     lesser scaup, 231<sup>7</sup>  
     ring-necked, 233<sup>1</sup>  
     ruddy, 239<sup>1</sup>  
     rufous-crested, 229<sup>7</sup>  
     wood, 229<sup>5</sup>  
 Dunlin, 265<sup>1</sup>  
**Eagle**, bald, 291<sup>1</sup>  
     golden, 289<sup>7</sup>  
*Ectopistes migratorius*, 283<sup>3</sup>  
 Egret, American, 247<sup>3</sup>  
 Eider, American, 235<sup>7</sup>  
     king, 237<sup>1</sup>  
*Elanoides forficatus*, 285<sup>3</sup>  
*Empidonax flaviventris*, 311<sup>5</sup>  
     *minimus*, 313<sup>3</sup>  
     *traillii alnorum*, 313<sup>1</sup>  
     *virescens*, 311<sup>7</sup>  
*Ereunetes occidentalis*, 267<sup>1</sup>  
     *pusillus*, 265<sup>7</sup>  
*Erismatura jamaicensis*, 239<sup>1</sup>  
**Falco** *columbarius*, 293<sup>3</sup>  
     *islandus*, 291<sup>3</sup>  
     *peregrinus anatum*, 293<sup>1</sup>  
     *rusticolus gyrfalco*, 291<sup>5</sup>  
         *obsoletus*, 291<sup>7</sup>  
     *sparverius*, 293<sup>5</sup>  
 Falconidae, 285<sup>3-95</sup><sup>3</sup>  
 Finch, purple, 323<sup>1</sup>  
 Flicker, northern, 305<sup>7</sup>

- Flycatcher, alder, 313<sup>1</sup>  
     crested, 309<sup>5</sup>  
     green-crested, 311<sup>7</sup>  
     least, 313<sup>3</sup>  
     olive-sided, 311<sup>1</sup>  
     scissor-tailed, 395<sup>3</sup>  
     yellow-bellied, 311<sup>5</sup>  
*Fratercula arctica*, 199<sup>5</sup>  
*Fregata aquila*, 223<sup>3</sup>  
*Fregatidae*, 223<sup>3</sup>  
*Fringillidae*, 321<sup>5</sup>-41<sup>6</sup>, 387<sup>5</sup>  
*Fulica americana*, 255<sup>7</sup>  
*Fulmar*, 389<sup>8</sup>  
     lesser, 391<sup>1</sup>  
*Fulmarus glacialis*, 389<sup>8</sup>  
     minor, 391<sup>1</sup>  
**Gadwall**, 225<sup>7</sup>  
*Galeoscoptes carolinensis*, 373<sup>3</sup>  
*Gallinae*, 279<sup>5</sup>-83<sup>3</sup>  
*Gallinago delicata*, 259<sup>5</sup>  
*Gallinula galeata*, 255<sup>5</sup>  
*Gallinule*, Florida, 255<sup>5</sup>  
     purple, 255<sup>3</sup>  
*Gannet*, 221<sup>1</sup>  
*Gavia arctica*, 199<sup>1</sup>  
     imber, 197<sup>8</sup>  
     lumme, 199<sup>3</sup>  
*Gaviidae*, 197<sup>8</sup>-99<sup>5</sup>  
*Gelochelidon nilotica*, 209<sup>7</sup>  
*Geothlypis agilis*, 367<sup>7</sup>  
     formosa, 367<sup>8</sup>  
     philadelphia, 369<sup>1</sup>  
     trichas, 369<sup>3</sup>  
*Gnatcatcher*, blue-gray, 381<sup>3</sup>  
*Godwit*, Hudsonian, 267<sup>7</sup>  
     marbled, 267<sup>3</sup>  
*Golden-eye*, American, 233<sup>3</sup>  
     Barrow's, 233<sup>5</sup>  
*Goldfinch*, American, 325<sup>3</sup>  
     European, 387<sup>5</sup>  
*Goose*, American white-fronted, 239<sup>7</sup>  
     barnacle, 243<sup>1</sup>  
     blue, 239<sup>5</sup>  
     Canada, 241<sup>1</sup>  
*Goose*, greater snow, 239<sup>3</sup>  
     Hutchins's, 241<sup>3</sup>  
*Goshawk*, American, 287<sup>8</sup>  
*Grackle*, bronzed, 321<sup>3</sup>  
     purple, 321<sup>1</sup>  
*Grebe*, Holboell's, 197<sup>8</sup>  
     horned, 197<sup>4</sup>  
     pied-billed, 197<sup>6</sup>  
*Grosbeak*, blue, 339<sup>5</sup>  
     evening, 321<sup>5</sup>  
     pine, 321<sup>7</sup>  
     rose-breasted, 339<sup>3</sup>  
*Grouse*, Canada, 279<sup>8</sup>  
     Canadian ruffed, 281<sup>3</sup>  
     ruffed, 281<sup>1</sup>  
*Gruidae*, 251<sup>1</sup>  
*Grus americana*, 251<sup>2</sup>  
     mexicana, 251<sup>2</sup>  
*Guara alba*, 243<sup>5</sup>  
*Guillemot*, black, 199<sup>7</sup>  
     Mandt's, 389<sup>2</sup>  
*Guiraca caerulea*, 339<sup>5</sup>  
*Gull*, American herring, 207<sup>3</sup>  
     Bonaparte's, 209<sup>1</sup>  
     glaucous, 205<sup>3</sup>  
     great black-backed, 205<sup>7</sup>  
     herring, 207<sup>1</sup>  
     Iceland, 205<sup>5</sup>  
     ivory, 203<sup>7</sup>  
     laughing, 207<sup>7</sup>  
     little, 209<sup>8</sup>  
     ring-billed, 207<sup>5</sup>  
     Sabine's, 209<sup>5</sup>  
*Gyr Falcon*, 291<sup>5</sup>  
     black, 291<sup>7</sup>  
     white, 291<sup>3</sup>  
**Haematopodidae**, 279<sup>8</sup>  
*Haematopus palliatus*, 279<sup>8</sup>  
*Haliaeetus leucocephalus*, 291<sup>1</sup>  
*Harelda hyemalis*, 235<sup>1</sup>  
*Harporhynchus rufus*, 373<sup>5</sup>  
*Hawk*, American rough-legged, 289<sup>5</sup>  
     American sparrow, 293<sup>3</sup>  
     broad-winged, 289<sup>3</sup>  
     Cooper's, 287<sup>1</sup>

Hawk, duck, 293<sup>1</sup>  
     marsh, 285<sup>5</sup>  
     pigeon, 293<sup>8</sup>  
     red-shouldered, 287<sup>7</sup>  
     red-tailed, 287<sup>5</sup>  
     sharp-shinned, 285<sup>7</sup>  
     Swainson's, 289<sup>1</sup>  
 Helminthophila celata, 357<sup>1</sup>  
     chrysoptera, 355<sup>1</sup>  
     lawrencei, 355<sup>8</sup>  
     leucobronchialis, 355<sup>5</sup>  
     peregrina, 357<sup>8</sup>  
     pinus, 353<sup>7</sup>  
     rubricapilla, 355<sup>7</sup>  
 Helmitherus vermivorus, 353<sup>5</sup>  
 Helodromas solitarius, 269<sup>5</sup>  
 Hen, heath, 281<sup>7</sup>  
 Herodiones, 243<sup>5-49</sup>  
 Heron, black-crowned night, 249<sup>8</sup>  
     great blue, 247<sup>1</sup>  
     green, 249<sup>8</sup>  
     little blue, 249<sup>1</sup>  
     Louisiana, 247<sup>7</sup>  
     snowy, 247<sup>5</sup>  
     yellow-crowned night, 249<sup>7</sup>  
 Hesperocichla naevia, 385<sup>8</sup>  
 Himantopus mexicanus, 259<sup>1</sup>  
 Hirundinidae, 343<sup>5-45</sup>  
 Hirundo erythrogaster, 345<sup>1</sup>  
 Histrionicus histrionicus, 235<sup>8</sup>  
 Hummingbird, ruby-throated, 307<sup>8</sup>  
 Hydrochelidon nigra surinamensis,  
     215<sup>8</sup>  
 Hylocichla aliciae, 383<sup>1</sup>  
     bicknelli, 383<sup>8</sup>  
     aonalaschkae pallasii, 383<sup>7</sup>  
     fuscescens, 381<sup>7</sup>  
     mustelina, 381<sup>5</sup>  
     ustulata swainsonii, 383<sup>5</sup>  
**Ibididae**, 243<sup>5-45</sup>  
 Ibis, glossy, 243<sup>7</sup>  
     white, 243<sup>5</sup>  
     white-faced glossy, 245<sup>1</sup>  
     wood, 245<sup>8</sup>  
 Icteria virens, 369<sup>8</sup>

Icteridae, 317<sup>8-21</sup>  
 Icterus galbula, 319<sup>5</sup>  
     spurius, 319<sup>3</sup>  
 Ionornis martinica, 255<sup>3</sup>  
**Jaeger**, long-tailed, 203<sup>8</sup>  
     parasitic, 203<sup>8</sup>  
     pomarine, 203<sup>1</sup>  
 Jay, blue, 315<sup>1</sup>  
     Canada, 315<sup>8</sup>  
 Junco, slate-colored, 335<sup>5</sup>  
 Junco hyemalis, 335<sup>5</sup>  
**Killdeer**, 275<sup>7</sup>  
 Kingbird, 309<sup>3</sup>  
     Arkansas, 309<sup>4</sup>  
 Kingfisher, belted, 301<sup>5</sup>  
 Kinglet, golden-crowned, 379<sup>7</sup>  
     ruby-crowned, 381<sup>1</sup>  
 Kite, swallow-tailed, 285<sup>3</sup>  
 Kittiwake, 205<sup>1</sup>  
 Knot, 261<sup>5</sup>  
**Lagopus lagopus**, 281<sup>5</sup>  
 Laniidae, 347<sup>5-49</sup>  
 Lanius borealis, 347<sup>5</sup>  
     ludovicianus, 347<sup>7</sup>  
     excubitorides, 349<sup>1</sup>  
 Lapwing, 275<sup>1</sup>  
 Laridae, 203<sup>7-15</sup>  
 Lark, horned, 313<sup>5</sup>  
     prairie horned, 313<sup>7</sup>  
 Larus argentatus, 207<sup>1</sup>  
     smithsonianus, 207<sup>8</sup>  
     atricilla, 207<sup>7</sup>  
     delawarensis, 207<sup>5</sup>  
     glaucus, 205<sup>8</sup>  
     leucopterus, 205<sup>5</sup>  
     marinus, 205<sup>7</sup>  
     minutus, 209<sup>8</sup>  
     philadelphia, 209<sup>1</sup>  
 Limicolae, 257<sup>1-79</sup>  
 Limosa fedoa, 267<sup>5</sup>  
     haemastica, 267<sup>7</sup>  
 Longipennes, 201<sup>7-15</sup>  
 Longspur, chestnut-collared, 327<sup>8</sup>  
     Lapland, 327<sup>1</sup>

- Loon, 197<sup>s</sup>  
     black-throated, 199<sup>1</sup>  
     red-throated, 199<sup>3</sup>  
 Lophodytes cucullatus, 225<sup>1</sup>  
 Loxia curvirostra minor, 323<sup>3</sup>  
     leucoptera, 323<sup>5</sup>  
**Machrochires**, 307<sup>1</sup>  
 Macrorhamphus griseus, 259<sup>7</sup>  
     scolopaceus, 261<sup>1</sup>  
 Magpie, American, 395<sup>5</sup>  
 Mallard, 225<sup>3</sup>  
 Man-o'-war bird, 223<sup>3</sup>  
 Mareca americana, 227<sup>8</sup>  
     penelope, 227<sup>1</sup>  
 Martin, purple, 343<sup>5</sup>  
 Meadowlark, 319<sup>1</sup>  
 Megalestris skua, 201<sup>7</sup>  
 Megascops asio, 297<sup>5</sup>  
 Melanerpes carolinus, 305<sup>5</sup>  
     erythrocephalus, 305<sup>3</sup>  
 Meleagris gallopavo fera, 283<sup>1</sup>  
 Melospiza fasciata, 335<sup>7</sup>  
     georgiana, 337<sup>8</sup>  
     lincolni, 337<sup>1</sup>  
 Merganser, American, 223<sup>5</sup>  
     hooded, 225<sup>1</sup>  
     red-breasted, 223<sup>7</sup>  
 Merganser americanus, 223<sup>6</sup>  
     serrator, 223<sup>7</sup>  
 Merula migratoria, 385<sup>1</sup>  
 Micropalma himantopus, 261<sup>8</sup>  
 Micropodidae, 307<sup>5</sup>  
 Milvulus forficatus, 395<sup>5</sup>  
 Mimus polyglottos, 373<sup>1</sup>  
 Mniotilta varia, 353<sup>1</sup>  
 Mniotiltidae, 353<sup>1</sup>-71<sup>7</sup>  
 Mockingbird, 373<sup>1</sup>  
 Molothrus ater, 317<sup>5</sup>  
 Motacillidae, 371<sup>7</sup>  
 Murre, 389<sup>4</sup>  
     Brünnich's, 201<sup>1</sup>  
 Myiarchus crinitus, 309<sup>5</sup>  
**Netta** rufina, 229<sup>7</sup>  
 Nettion carolinensis, 227<sup>8</sup>  
     crecca, 391<sup>5</sup>  
 Nighthawk, 307<sup>3</sup>  
 Numenius arquatus, 273<sup>7</sup>  
     borealis, 273<sup>5</sup>  
     hudsonicus, 273<sup>8</sup>  
     longirostris, 273<sup>1</sup>  
 Nuthatch, brown-headed, 377<sup>7</sup>  
     red-breasted, 377<sup>5</sup>  
     white-breasted, 377<sup>8</sup>  
 Nyctala acadica, 297<sup>8</sup>  
     tengmalmi richardsoni, 393<sup>7</sup>  
 Nyctea nyctea, 299<sup>1</sup>  
 Nycticorax nycticorax naevius, 249<sup>5</sup>  
     violaceus, 249<sup>7</sup>  
**Oceanites** oceanicus, 219<sup>4</sup>  
 Oceanodroma leucorhoa, 219<sup>8</sup>  
 Oidemia americana, 237<sup>8</sup>  
     deglandi, 237<sup>5</sup>  
     perspicillata, 237<sup>7</sup>  
 Old-squaw, 235<sup>1</sup>  
 Olor columbianus, 243<sup>3</sup>  
 Oriole, Baltimore, 319<sup>5</sup>  
     orchard, 319<sup>3</sup>  
 Osprey, American, 293<sup>7</sup>  
 Otocoris alpestris, 313<sup>5</sup>  
     praticola, 313<sup>7</sup>  
 Oven-bird, 365<sup>7</sup>  
 Owl, American barn, 295<sup>1</sup>  
     American hawk, 299<sup>3</sup>  
     American long-eared, 295<sup>8</sup>  
     barred, 295<sup>7</sup>  
     burrowing, 299<sup>5</sup>  
     great gray, 297<sup>1</sup>  
     great horned, 297<sup>7</sup>  
     Richardson's, 393<sup>7</sup>  
     saw-whet, 297<sup>8</sup>  
     screech, 297<sup>5</sup>  
     short-eared, 295<sup>5</sup>  
     snowy, 299<sup>1</sup>  
 Oyster-catcher, American, 279<sup>8</sup>  
**Pagophila** alba, 203<sup>7</sup>  
 Paludicola, 251<sup>1</sup>-55<sup>9</sup>  
 Pandion haliaëtus carolinensis, 293<sup>7</sup>  
 Paridae, 377<sup>8</sup>-79<sup>7</sup>  
 Paroquet, Carolina, 299<sup>7</sup>  
 Parrots, 299<sup>7</sup>



- Parus atricapillus*, 379<sup>3</sup>  
     *bicolor*, 379<sup>1</sup>  
     *carolinensis*, 397<sup>1</sup>  
     *hudsonicus*, 379<sup>5</sup>  
*Passer domesticus*, 387<sup>7</sup>  
*Passerella iliaca*, 337<sup>5</sup>  
*Passeres*, 309<sup>1-85</sup>  
*Passerina nivalis*, 325<sup>7</sup>  
*Pavoncella pugnax*, 271<sup>1</sup>  
*Pelecanidae*, 223<sup>1</sup>  
*Pelecanus erythrorhynchos*, 223<sup>1</sup>  
     *fuscus*, 391<sup>3</sup>  
 Pelican, American white, 223<sup>1</sup>  
     brown, 391<sup>3</sup>  
 Perching birds, 309<sup>1-87</sup>  
*Perisoreus canadensis*, 315<sup>3</sup>  
*Petrel*, black-capped, 217<sup>7</sup>  
     Leach's, 219<sup>3</sup>  
     scaled, 219<sup>1</sup>  
     Wilson's, 219<sup>4</sup>  
*Petrochelidon lunifrons*, 343<sup>7</sup>  
 Pewee, wood, 311<sup>3</sup>  
*Phaëthon americanus*, 219<sup>7</sup>  
*Phaëthontidae*, 219<sup>7</sup>  
*Phalacrocoracidae*, 221<sup>5</sup>  
*Phalacrocorax carbo*, 221<sup>5</sup>  
     *dilophus*, 221<sup>7</sup>  
*Phalarope*, northern, 257<sup>3</sup>  
     red, 257<sup>2</sup>  
     Wilson's, 257<sup>5</sup>  
*Phalaropodidae*, 257<sup>1</sup>  
*Phalaropus lobatus*, 257<sup>3</sup>  
*Phasianidae*, 283<sup>1</sup>  
*Philohela minor*, 259<sup>3</sup>  
 Phoebe, 309<sup>7</sup>  
*Pica pica hudsonica*, 395<sup>5</sup>  
*Pici*, 301<sup>7-5</sup>  
*Picidae*, 301<sup>7-5</sup>  
*Picoides americanus*, 303<sup>5</sup>  
     *arcticus*, 303<sup>3</sup>  
 Pigeon, passenger, 283<sup>3</sup>  
*Pinicola enucleator canadensis*, 321<sup>7</sup>  
*Pintail*, 229<sup>3</sup>  
*Pipilo erythrophthalmus*, 337<sup>7</sup>  
*Pipit*, American, 371<sup>7</sup>  
*Piranga erythromelas*, 343<sup>1</sup>  
     *ludoviciana*, 341<sup>7</sup>  
     *rubra*, 343<sup>3</sup>  
*Plegadis autumnalis*, 243<sup>7</sup>  
     *guarauna*, 245<sup>1</sup>  
*Plover*, American golden, 275<sup>5</sup>  
     belted piping, 277<sup>5</sup>  
     black-bellied, 275<sup>3</sup>  
     piping, 277<sup>3</sup>  
     semipalmated, 277<sup>1</sup>  
     Wilson's, 277<sup>7</sup>  
*Podicipidae*, 197<sup>3</sup>  
*Podilymbus podiceps*, 197<sup>5</sup>  
*Polioptila caerulea*, 381<sup>3</sup>  
*Poocetes gramineus*, 327<sup>5</sup>  
*Porzana carolina*, 253<sup>3</sup>  
     *jamaicensis*, 253<sup>7</sup>  
     *noveboracensis*, 253<sup>5</sup>  
*Procellariidae*, 215<sup>7-19</sup>  
*Progne subis*, 343<sup>5</sup>  
*Protonotaria citrea*, 353<sup>1</sup>  
*Psittaci*, 299<sup>7</sup>  
*Psittacidae*, 299<sup>7</sup>  
*Ptarmigan*, willow, 281<sup>1</sup>  
*Puffin*, 199<sup>5</sup>  
*Puffinus auduboni*, 217<sup>3</sup>  
     *borealis*, 215<sup>7</sup>  
     *fuliginosus*, 217<sup>5</sup>  
     *gravis*, 217<sup>1</sup>  
*Pygopodes*, 197<sup>3-201</sup>  
*Querquedula cyanoptera*, 391<sup>3</sup>  
     *discors*, 227<sup>7</sup>  
*Quiscalus quiscula*, 321<sup>1</sup>  
     *aeneus*, 321<sup>3</sup>  
**Rail**, black, 253<sup>7</sup>  
     clapper, 251<sup>7</sup>  
     king, 251<sup>5</sup>  
     Virginia, 253<sup>1</sup>  
     yellow, 253<sup>5</sup>  
*Rallidae*, 251<sup>3-55</sup>  
*Rallus crepitans*, 251<sup>7</sup>  
     *elegans*, 251<sup>5</sup>  
     *virginianus*, 253<sup>1</sup>  
*Raptores*, 283<sup>7-99</sup>  
*Raven*, northern, 315<sup>3</sup>

- Recurvirostra americana*, 257<sup>a</sup>  
*Recurvirostridae*, 257<sup>a</sup>-59<sup>a</sup>  
 Redhead, 231<sup>1</sup>  
 Redpoll, 323<sup>7</sup>  
     greater, 325<sup>1</sup>  
 Redstart, American, 371<sup>5</sup>  
*Regulus calendula*, 381<sup>1</sup>  
     satrapa, 379<sup>7</sup>  
*Rissa tridactyla*, 205<sup>1</sup>  
 Robin, American, 385<sup>1</sup>  
 Ruff, 271<sup>1</sup>  
*Rynchopidae*, 215<sup>5</sup>  
*Rynchops nigra*, 215<sup>5</sup>  
**Sanderling**, 267<sup>3</sup>  
*Sandpiper*, Baird's, 263<sup>5</sup>  
     Bartramian, 271<sup>3</sup>  
     buff-breasted, 271<sup>5</sup>  
     curlew, 265<sup>5</sup>  
     least, 263<sup>7</sup>  
     pectoral, 263<sup>1</sup>  
     purple, 261<sup>7</sup>  
     red-backed, 265<sup>3</sup>  
     semipalmated, 265<sup>7</sup>  
     solitary, 269<sup>5</sup>  
     spotted, 271<sup>7</sup>  
     stilt, 261<sup>3</sup>  
     western, 267<sup>1</sup>  
     white-rumped, 263<sup>3</sup>  
*Sapsucker*, yellow-bellied, 303<sup>7</sup>  
*Saxicola oenanthe*, 385<sup>5</sup>  
*Sayornis phoebe*, 309<sup>7</sup>  
*Scolecophagus carolinus*, 319<sup>7</sup>  
*Scolopacidae*, 259<sup>a</sup>-73<sup>a</sup>  
*Scolopax rusticola*, 393<sup>5</sup>  
 Scoter, American, 237<sup>1</sup>  
     surf, 237<sup>7</sup>  
     white-winged, 237<sup>5</sup>  
*Scotiaptex cinerea*, 297<sup>1</sup>  
*Seiurus aurocapillus*, 365<sup>7</sup>  
     motacilla, 367<sup>a</sup>  
     noveboracensis, 367<sup>1</sup>  
*Setophaga ruticilla*, 371<sup>5</sup>  
*Shearwater*, Audubon's, 217<sup>3</sup>  
     Cory's, 215<sup>7</sup>  
*Shearwater*, greater, 217<sup>1</sup>  
     sooty, 217<sup>5</sup>  
*Shore birds*, 257<sup>1</sup>-79<sup>5</sup>  
 Shoveller, 229<sup>1</sup>  
 Shrike, loggerhead, 347<sup>7</sup>  
     northern, 347<sup>5</sup>  
     white-rumped, 349<sup>1</sup>  
*Sialia sialis*, 385<sup>7</sup>  
 Siskin, pine, 325<sup>5</sup>  
*Sitta canadensis*, 377<sup>5</sup>  
     carolinensis, 377<sup>3</sup>  
     pusilla, 377<sup>7</sup>  
 Skimmer, black, 215<sup>5</sup>  
 Skua, 201<sup>7</sup>  
 Skylark, 387<sup>1</sup>  
 Snipe, Wilson's, 259<sup>5</sup>  
 Snowflake, 325<sup>7</sup>  
*Somateria dresseri*, 235<sup>7</sup>  
     spectabilis, 237<sup>1</sup>  
 Sora, 253<sup>3</sup>  
*Sparrow*, Acadian sharp-tailed, 331<sup>5</sup>  
     chipping, 335<sup>1</sup>  
     field, 335<sup>3</sup>  
     fox, 337<sup>5</sup>  
     grasshopper, 329<sup>3</sup>  
     Henslow's, 329<sup>5</sup>  
     house, 387<sup>7</sup>  
     Ipswich, 327<sup>7</sup>  
     lark, 333<sup>1</sup>  
     Leconte's, 329<sup>7</sup>  
     Lincoln's, 337<sup>1</sup>  
     Nelson's, 331<sup>3</sup>  
     Savanna, 329<sup>1</sup>  
     seaside, 331<sup>7</sup>  
     sharp-tailed, 331<sup>1</sup>  
     song, 335<sup>7</sup>  
     swamp, 337<sup>3</sup>  
     tree, 333<sup>7</sup>  
     vesper, 327<sup>5</sup>  
     white-crowned, 333<sup>3</sup>  
     white-throated, 333<sup>5</sup>  
*Spatula clypeata*, 229<sup>1</sup>  
*Speotyto cunicularia hypogaea*, 299<sup>5</sup>  
*Sphyrapicus varius*, 303<sup>7</sup>  
*Spinus pinus*, 325<sup>5</sup>  
*Spiza americana*, 341<sup>3</sup>

- Spizella monticola*, 333<sup>r</sup>  
     *pusilla*, 335<sup>s</sup>  
     *socialis*, 335<sup>r</sup>  
*Squatarola squatarola*, 275<sup>s</sup>  
 Starling, 387<sup>s</sup>  
*Steganopodes*, 219<sup>s</sup>-23<sup>s</sup>  
*Steganopus tricolor*, 257<sup>s</sup>  
*Stelgidopteryx serripennis*, 345<sup>r</sup>  
*Stercorariidae*, 201<sup>r</sup>-3<sup>r</sup>  
*Stercorarius longicaudus*, 203<sup>s</sup>  
     *parasiticus*, 203<sup>s</sup>  
     *pomarinus*, 203<sup>r</sup>  
*Sterna antillarum*, 213<sup>r</sup>  
     *caspia*, 211<sup>r</sup>  
     *dougalli*, 213<sup>s</sup>  
     *forsteri*, 211<sup>r</sup>  
     *fuliginosa*, 215<sup>r</sup>  
     *hirundo*, 213<sup>r</sup>  
     *maxima*, 211<sup>s</sup>  
     *paradisaea*, 213<sup>s</sup>  
     *sandvicensis acuflava*, 389<sup>s</sup>  
     *trudeaui*, 211<sup>s</sup>  
 Stilt, black-necked, 259<sup>r</sup>  
*Strigidae*, 295<sup>r</sup>  
*Strix pratincola*, 295<sup>r</sup>  
*Sturnella magna*, 319<sup>r</sup>  
*Sturnidae*, 387<sup>s</sup>  
*Sturnus vulgaris*, 387<sup>s</sup>  
*Sula bassana*, 221<sup>s</sup>  
     *sula*, 221<sup>r</sup>  
*Sulidae*, 221<sup>r</sup>  
 Surf birds, 279<sup>r</sup>  
*Surnia ulula caparoch*, 299<sup>s</sup>  
 Swallow, bank, 345<sup>s</sup>  
     barn, 345<sup>r</sup>  
     cliff, 343<sup>r</sup>  
     rough-winged, 345<sup>r</sup>  
     tree, 345<sup>s</sup>  
 Swan, whistling, 243<sup>s</sup>  
 Swift, chimney, 307<sup>s</sup>  
*Swimmers, lamellirostral*, 223<sup>s</sup>-43<sup>s</sup>  
     long-winged, 201<sup>r</sup>-15<sup>s</sup>  
     totipalmate, 219<sup>s</sup>-23<sup>s</sup>  
     tube-nosed, 215<sup>r</sup>-19<sup>s</sup>  
*Sylviidae*, 379<sup>r</sup>-81<sup>s</sup>  
*Symphemia semipalmata*, 269<sup>r</sup>  
*Syrnium nebulosum*, 295<sup>r</sup>  
*Tachycineta bicolor*, 345<sup>s</sup>  
 Tanager, Louisiana, 341<sup>r</sup>  
     scarlet, 343<sup>r</sup>  
     summer, 343<sup>s</sup>  
*Tanagridae*, 341<sup>r</sup>-43<sup>s</sup>  
*Tantalus loculator*, 245<sup>s</sup>  
 Teal, blue-winged, 227<sup>r</sup>  
     cinnamon, 391<sup>s</sup>  
     European, 391<sup>s</sup>  
     green-winged, 227<sup>s</sup>  
 Tern, arctic, 213<sup>s</sup>  
     black, 215<sup>s</sup>  
     Cabot's, 389<sup>s</sup>  
     Caspian, 211<sup>r</sup>  
     common, 213<sup>r</sup>  
     Forster's, 211<sup>r</sup>  
     gull-billed, 209<sup>r</sup>  
     least, 213<sup>r</sup>  
     roseate, 213<sup>s</sup>  
     royal, 211<sup>s</sup>  
     sooty, 215<sup>r</sup>  
     Trudeau's, 211<sup>s</sup>  
*Tetraonidae*, 279<sup>s</sup>-81<sup>s</sup>  
 Thrasher, brown, 373<sup>s</sup>  
 Thrush, Bicknell's, 383<sup>s</sup>  
     gray-cheeked, 383<sup>r</sup>  
     hermit, 383<sup>r</sup>  
     olive-backed, 383<sup>s</sup>  
     varied, 385<sup>s</sup>  
     Wilson's, 381<sup>r</sup>  
     wood, 381<sup>s</sup>  
*Thryothorus ludovicianus*, 373<sup>r</sup>  
 Titmouse, tufted, 379<sup>r</sup>  
*Totanus flavipes*, 269<sup>s</sup>  
     *melanoleucus*, 269<sup>r</sup>  
 Towhee, 337<sup>r</sup>  
*Tringa alpina*, 265<sup>r</sup>  
     *pacifica*, 265<sup>s</sup>  
     *bairdii*, 263<sup>s</sup>  
     *canutus*, 261<sup>s</sup>  
     *ferruginea*, 265<sup>s</sup>  
     *fuscescens*, 263<sup>s</sup>  
     *maculata*, 263<sup>r</sup>  
     *maritima*, 261<sup>r</sup>  
     *minutilla*, 263<sup>r</sup>  
*Trochilidae*, 307<sup>r</sup>

- Trochilus colubris*, 307<sup>8</sup>  
*Troglodytes aëdon*, 375<sup>1</sup>  
*Troglodytidae*, 373<sup>1</sup>-75<sup>9</sup>  
 Tropic bird, yellow-billed, 219<sup>7</sup>  
*Tryngites subruficollis*, 271<sup>5</sup>  
*Tubinares*, 215<sup>7</sup>-19<sup>9</sup>  
*Turdidae*, 381<sup>5</sup>-85<sup>9</sup>  
 Turkey, wild, 283<sup>1</sup>  
 Turnstone, 279<sup>1</sup>  
*Tympanuchus cupido*, 281<sup>7</sup>  
*Tyrannidae*, 309<sup>1</sup>-13<sup>5</sup>  
*Tyrannus tyrannus*, 309<sup>2</sup>  
     *verticalis*, 309<sup>4</sup>  
**Uria lomvia**, 201<sup>1</sup>  
     *troile*, 389<sup>4</sup>  
**Vanellus vanellus**, 275<sup>1</sup>  
*Vireo*, blue-headed, 351<sup>3</sup>  
     *Philadelphia*, 349<sup>8</sup>  
     *plumbeus*, 351<sup>5</sup>  
     *red-eyed*, 349<sup>8</sup>  
     *warbling*, 349<sup>7</sup>  
     *white-eyed*, 351<sup>7</sup>  
     *yellow-throated*, 351<sup>1</sup>  
*Vireo flavifrons*, 351<sup>1</sup>  
     *gilvus*, 349<sup>7</sup>  
     *noveboracensis*, 351<sup>7</sup>  
     *olivaceus*, 349<sup>3</sup>  
     *philadelphicus*, 349<sup>5</sup>  
     *solitarius*, 351<sup>3</sup>  
         *plumbeus*, 351<sup>5</sup>  
*Vireonidae*, 349<sup>3</sup>-51<sup>9</sup>  
*Vulture*, black, 285<sup>1</sup>  
     *turkey*, 283<sup>7</sup>  
**Wagtails**, 371<sup>7</sup>  
**Warbler**, bay-breasted, 361<sup>5</sup>  
     black and white, 353<sup>1</sup>  
     Blackburnian, 363<sup>1</sup>  
     black poll, 361<sup>7</sup>  
     black-throated blue, 359<sup>3</sup>  
     black-throated green, 363<sup>5</sup>  
     blue-winged, 353<sup>7</sup>  
     Brewster's, 355<sup>5</sup>  
     Canadian, 371<sup>8</sup>  
     Cape May, 357<sup>7</sup>  
     *Warbler*, cerulean, 361<sup>1</sup>  
         chestnut-sided, 361<sup>3</sup>  
         Connecticut, 367<sup>7</sup>  
         golden-winged, 355<sup>1</sup>  
         hooded, 369<sup>7</sup>  
         Kentucky, 367<sup>5</sup>  
         Lawrence's, 355<sup>3</sup>  
         magnolia, 359<sup>9</sup>  
         mourning, 369<sup>1</sup>  
         myrtle, 359<sup>5</sup>  
         Nashville, 355<sup>7</sup>  
         northern parula, 357<sup>5</sup>  
         orange-crowned, 357<sup>1</sup>  
         palm, 365<sup>1</sup>  
         pine, 363<sup>7</sup>  
         prairie, 365<sup>5</sup>  
         prothonotary, 353<sup>9</sup>  
         Tennessee, 357<sup>3</sup>  
         Wilson's, 371<sup>1</sup>  
         worm-eating, 353<sup>5</sup>  
         yellow, 359<sup>1</sup>  
         yellow palm, 365<sup>3</sup>  
         yellow-throated, 363<sup>3</sup>  
 Water-thrush, 367<sup>1</sup>  
     Louisiana, 367<sup>3</sup>  
 Waxwing, Bohemian, 347<sup>1</sup>  
     cedar, 347<sup>3</sup>  
 Wheatear, 385<sup>5</sup>  
 Whip-poor-will, 307<sup>2</sup>  
 Widgeon, 227<sup>1</sup>  
 Willet, 269<sup>7</sup>  
*Wilsonia canadensis*, 371<sup>8</sup>  
     *mitrata*, 369<sup>7</sup>  
     *pusilla*, 371<sup>1</sup>  
 Wood warblers, 353<sup>1</sup>-71<sup>7</sup>  
 Woodcock, American, 259<sup>3</sup>  
     European, 393<sup>5</sup>  
 Woodpecker, American three-toed,  
     303<sup>5</sup>  
     arctic three-toed, 303<sup>3</sup>  
     downy, 303<sup>1</sup>  
     hairy, 301<sup>8</sup>  
     northern pileated, 305<sup>1</sup>  
     red-bellied, 305<sup>5</sup>  
     red-cockaded, 395<sup>1</sup>  
     red-headed, 305<sup>3</sup>

Wren, Carolina, 373<sup>7</sup>

house, 375<sup>1</sup>

long-billed marsh, 375<sup>7</sup>

short-billed marsh, 375<sup>5</sup>

winter, 375<sup>3</sup>

**Xanthocephalus** xanthocephalus,  
395<sup>7</sup>

Xema sabinii, 209<sup>5</sup>

**Yellow-legs**, 269<sup>3</sup>

greater, 269<sup>1</sup>

Yellow-throat, Maryland, 369<sup>3</sup>

**Zamelodia ludoviciana**, 339<sup>1</sup>

Zenaidura macroura, 283<sup>1</sup>

Zonotrichia albicollis, 333<sup>5</sup>

leucophrys, 333<sup>3</sup>



New York State Museum

PUBLICATIONS

**Museum reports.** New York state museum. Annual report, 1847–date. O. Albany 1848–date.

Average 300 pages a year. Price for all in print to 1892, 50 cents a volume; 75 cents in cloth; 1892–date, 75 cents, cloth.

**Museum bulletins.** New York state museum. Bulletins. v. 1–6, O. Albany 1887–date. Price to advance subscribers, 75 cents a year.

**Volume 1.** 6 nos. Price \$1 in cloth

- 1 Marshall, W: B. Preliminary list of New York unionidae. 19p. Mar. 1892. Price 5 cents.
- 2 Peck, C: H. Contributions to the botany of the state of New York. 66p. 2 pl. May 1887. Price [35] cents.
- 3 Smock, J: C. Building stone in the state of New York. 152p. Mar. 1888. Out of print.
- 4 Nason, F. L. Some New York minerals and their localities. 19p. 1 pl. Aug. 1888. Price 5 cents.
- 5 Lintner, J. A. White grub of the May beetle. 31p. il. Nov. 1888. Price 10 cents.
- 6 — Cut-worms. 36p. il. Nov. 1888. Price 10 cents.

**Volume 2.** 4 nos. Price \$1 in cloth

- 7 Smock, J: C. First report on the iron mines and iron ore districts in N. Y. 5+70p. map 58×60 cm. June 1889. Price 20 cents.
- 8 Peck, C: H. Boleti of the U. S. 96p. Sep. 1889. Price [40] cents.
- 9 Marshall, W: B. Beaks of unionidae inhabiting the vicinity of Albany, N. Y. 23p. 1 pl. Aug. 1890. Price 10 cents.
- 10 Smock, J: C. Building stone in New York. 210p. map 58×60 cm, tab. Sep. 1890. Price 40 cents.

**Volume 3**

- 11 Merrill, F: J. H. Salt and gypsum industries in New York. 92p. 12 pl. 2 maps 38×58, 61×66 cm, 11 tab. Ap. 1893. Price 40 cents.
- 12 Merrill, F: J. H. & Ries, Heinrich. Clay industries of New York. 174p. 2 pl. map 59×67 cm. Mar. 1895. Price 30 cents.  
New edition in preparation.
- 13 Lintner, J. A. Some destructive insects of New York state; San José scale. 54p. 7 pl. Ap. 1895. Price 15 cents.
- 14 Kemp, J. F. Geology of Moriah and Westport townships, Essex co. N. Y., with notes on the iron mines. 38p. 7 pl. 2 maps 30×33, 38×44 cm. Sep. 1895. Price 10 cents.
- 15 Merrill, F: J. H. Mineral resources of New York. 224p. 2 maps 22×35, 58×65 cm. Feb. 1896. Price 40 cents.

**Volume 4**

- 16 Beauchamp, W: M. Aboriginal chipped stone implements of New York. 86p. 23 pl. Oct. 1897. Price 25 cents.
- 17 Merrill, F: J. H. Road materials and road building in New York. 52p. 14 pl. 2 maps 34×44, 68×92 cm. Oct. 1897. Price 15 cents.
- 18 Beauchamp, W: M. Polished stone articles used by the New York aborigines. 104p. 35 pl. Nov. 1897. Price 25 cents.
- 19 Merrill, F: J. H. Guide to the study of the geological collections of the New York state museum. 162p. 119 pl. 1 map 33×43 cm. Nov. 1898. Price 40 cents.

**Volume 5**

- 20 Felt, E. P. Elm-leaf beetle in New York. 46p. il. 5 pl. June 1898. Price 5 cents.
- 21 Kemp, J. F. Geology of the Lake Placid region. 24p. 1 map 33×34 cm. 1 pl. Sep. 1898. Price 5 cents.

- 22 Beauchamp, W: M. Earthenware of the New York aborigines. 78p. 33 pl. Oct. 1898. *Price 25 cents.*
- 23 Felt, E. P. 14th report of the state entomologist 1898. 150p. il. 9 pl. Dec. 1898. *Price 20 cents.*
- 24 ——— Memorial of the life and entomologic work of J. A. Lintner. 316p. 1 pl. Oct. 1899. *Price 35 cents.*
- 25 Peck, C: H. Report of the state botanist 1898. 76p. 5 pl. Oct. 1899. *Price 40 cents.*

Volume 6

- 26 Felt, E. P. Collection, preservation and distribution of New York insects. 36p. il. Ap. 1899. *Price 5 cents.*
- 27 ——— Shade-tree pests in New York state. 26p. 5 pl. May 1899. *Price 5 cents.*
- 28 Peck, C: H. Plants of North Elba. 206p. 1 map 12×16 cm. June 1899. *Price 20 cents.*
- 29 Miller, G. S., jr. Preliminary list of New York mammals. 124p. Oct. 1899. *Price 15 cents.*
- 30 Orton, Edward. Petroleum and natural gas in New York. 136p. il. 3 maps 13×23, 7×22, 9×14 cm. Nov. 1899. *Price 15 cents.*
- 31 Felt, E. P. 15th report of the state entomologist 1899. *In preparation.*

Volume 7

- 32 Beauchamp, W: M. Aboriginal occupation of New York. 190p. 16 pl. 2 maps 44×35, 93×68 cm. March 1900. *Price 30 cents.*
  - 33 Farr, M. S. Check list of New York birds. 222p. Ap. 1900. *Price 25 cents.*
  - 34 Cumings, E. R. Lower Silurian system of eastern Montgomery county; Prosser, C: S. Notes on the stratigraphy of the Mohawk valley and Saratoga county, N. Y. *In press.*
  - Clarke, J: M. Paleontologic papers; Simpson, G: B. Preliminary descriptions of new genera. *In press.*
  - Ries, Heinrich. Clays: their properties and uses. *In press.*
  - Miller, G. S., jr. Key to the land mammals of the North America. *In press.*
  - Bean, Tarleton. Check list of the fishes of N. Y. *In preparation.*
- Economic map.** Merrill, F: J. H. Economic and geologic map of the state of New York. 59×67 cm. 1894. *Price, unmounted 25 cents, backed on muslin 75 cents.*  
Scale 14 miles to 1 inch. New edition in preparation.

**Museum memoirs.** New York state museum. Memoirs. v.1, Q. Albany 1889.

Uniform with the *Paleontology*.

- 1 Beecher, C: E. & Clarke, J: M. Development of some Silurian brachiopoda. 95p. 8 pl. Oct. 1889. *Price 80 cents.*
- 2 Hall, James & Clarke, J: M. Paleozoic reticulate sponges. 350p. il. 70 pl. Oct. 1899. *Price \$1.*
- 3 Clarke, J: M. The Oriskany fauna of Becraft mountain, Columbia co. N. Y. *In preparation.*

**Natural history.** New York state. Natural history of New York. 30 v. il. pl. maps, sq. Q. Albany 1842-94.

Divisions 1-5 out of print.


- Division 1** De Kay, J. E. Zoology. 5 v. 1842-44.  
" 2 Torrey, John. Botany. 2 v. 1843.  
" 3 Beck, L. C. Mineralogy. 24+536p. 1842.  
" 4 Mather, W: W.; Emmons, Ebenezer; Vanuxem, Lardner & Hall, James. Geology. 4 v. 1842-43.  
" 5 Emmons, Ebenezer. Agriculture. 5 v. 1846-54.  
" 6 Hall, James. Paleontology. 8 v. 1847-94.














SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01300 6820